



BlueCross BlueShield of Louisiana

An independent licensee of the Blue Cross and Blue Shield Association.

Fentanyl Oral Transmucosal and Nasal Opioid Analgesics

Policy # 00213

Original Effective Date: 12/20/2006

Current Effective Date: 06/18/2014

Applies to all products administered or underwritten by Blue Cross and Blue Shield of Louisiana and its subsidiary, HMO Louisiana, Inc. (collectively referred to as the "Company"), unless otherwise provided in the applicable contract. Medical technology is constantly evolving, and we reserve the right to review and update Medical Policy periodically.

When Services May Be Eligible for Coverage

Coverage for eligible medical treatments or procedures, drugs, devices or biological products may be provided only if:

- *Benefits are available in the member's contract/certificate, and*
- *Medical necessity criteria and guidelines are met.*

Based on review of available data, the Company may consider fentanyl oral or nasal opioid analgesics for the treatment of breakthrough cancer pain in patients with malignancies already receiving opioid therapy to be **eligible for coverage**.

Patient Selection Criteria

The use of fentanyl oral or nasal opioid analgesics will be considered for coverage eligibility when the following criteria are met:

- Patients with malignancies experiencing breakthrough cancer pain who are already receiving opioid therapy; and
- Patient must be opioid tolerant; and
- Requested drug is oral transmucosal fentanyl citrate (Actiq): Patient must be 16 years of age or older; and
- Requested drug is a fentanyl citrate buccal film or tablet (Onsolis, Fentora), sublingual product (Abstral, Subsys), or nasal spray (Lazanda): Patient must be 18 years of age or older.

When Services Are Considered Investigational

Coverage is not available for investigational medical treatments or procedures, drugs, devices or biological products.

Based on review of available data, the Company considers the use of fentanyl oral or nasal opioid analgesics when patient selection criteria are not met to be **investigational**.*

Background/Overview

Breakthrough pain characteristically has a rapid onset with moderate to severe intensity but brief duration. Pain of such intense severity but short duration is difficult to treat because currently available oral agents administered at the onset of a breakthrough pain episode may require 30-45 minutes to be absorbed and produce an analgesic effect.

Patients considered opioid tolerant are those who are taking at least 60mg of oral morphine /day, or at least 25mcg of transdermal fentanyl/hour, or at least 30mg of oxycodone daily, or at least 8mg of oral hydromorphone daily or an equianalgesic dose of another opioid for a week or longer.



BlueCross BlueShield of Louisiana

An independent licensee of the Blue Cross and Blue Shield Association.

Fentanyl Oral Transmucosal and Nasal Opioid Analgesics

Policy # 00213

Original Effective Date: 12/20/2006

Current Effective Date: 06/18/2014

Actiq delivers pain relief via a single-use medicated lozenge that is attached to a handle. It dissolves as the patient moves the unit along the inside of the cheek.

Fentanyl buccal tablet (Fentora) is a sugar-free tablet that is placed along the inside of the cheek. These medications utilize the oral transmucosal system to treat breakthrough cancer pain.

Onsolis (fentanyl buccal soluble film) is an oral transmucosal form of the potent opioid analgesic, fentanyl citrate, intended for application to the buccal mucosa. Onsolis uses the BioErodible MucoAdhesive (BEMA[™])[†] bilayer delivery technology which is comprised of water-soluble polymeric films. Onsolis consists of a pink bioadhesive layer bonded onto a white inactive layer. The active ingredient, fentanyl citrate, is incorporated into the bioadhesive layer, which adheres to the moist buccal mucosa. The amount of fentanyl delivered transmucosally is proportional to the film surface area. It is believed that the inactive layer isolates the bioadhesive layer from the saliva, which may optimize delivery of fentanyl across the buccal mucosa.

Subsys (fentanyl sublingual spray) is a potent opioid analgesic intended for sublingual mucosal administration. Subsys is formulated to be sprayed underneath the tongue to allow for absorption of fentanyl across the sublingual mucosa.

FDA or Other Governmental Regulatory Approval

U.S. Food and Drug Administration (FDA)

The FDA approves Actiq, Fentora, Onsolis, Lazanda, Abstral and Subsys for the management of breakthrough cancer pain in patients with malignancies who are already receiving and who are tolerant to opioid therapy. Because fentanyl is a schedule II narcotic, the FDA recommends that the product be packaged and marketed to minimize the opportunity for diversion, abuse or access to children.

Rationale/Source

The goals of cancer pain therapy are to find the right dose of around-the-clock medications to control persistent pain and the right dose of supplemental medication to relieve breakthrough pain. Treatment of breakthrough pain requires an effective analgesic that has a rapid onset of analgesia, a short duration of effect and a route of administration that is appropriate for the patient's condition.

Fentanyl is a pure opioid agonist whose principal therapeutic action is analgesia. The pharmacological effects of opioid agonists include anxiolysis, euphoria, feelings of relaxation, respiratory depression, constipation, miosis, cough suppression and analgesia.

Actiq, Fentora and Onsolis are intended to be used in the care of cancer patients and only by oncologists and pain specialists who are knowledgeable of and skilled in the use of schedule II opioids to treat cancer pain.

The absorption of the transmucosal dosage form is a combination of an initial rapid absorption from the buccal mucosa and a more prolonged absorption of swallowed fentanyl from the gastrointestinal tract. Approximately 25% of the total dose of transmucosal fentanyl is rapidly absorbed from the buccal mucosa



BlueCross BlueShield of Louisiana

An independent licensee of the Blue Cross and Blue Shield Association.

Fentanyl Oral Transmucosal and Nasal Opioid Analgesics

Policy # 00213

Original Effective Date: 12/20/2006

Current Effective Date: 06/18/2014

and becomes systemically available. The remaining 75% of the total dose is swallowed with the saliva and then is slowly absorbed from the gastrointestinal tract.

Studies on the relationship of age to the effects of transmucosal fentanyl have not been performed in the pediatric population. Transmucosal fentanyl contains a medicine in an amount that can be fatal to children. No difference was noted in the safety profile of the patients group over 65 years of age as compared to younger patients in clinical trials. The older population did titrate to a slightly lower dose.

References

1. ACTIQ.com. About Breakthrough Cancer Pain. 2004.
2. American Society of Health-System Pharmacists. AHFS Drug Information 2007. Fentanyl Citrate.
3. Fentora.com
4. www.Onsolis.com
5. Food and Drug Administration. Labeling of the Drug Actiq. September 2009.
6. Food and Drug Administration. Labeling of the Drug Fentora. December 5, 2009.
7. Food and Drug Administration. Labeling of the Drug Abstral. January 7, 2011.
8. Food and Drug Administration. Labeling of the Drug Subsys. January 1, 2012.
9. Medline Plus. Drug Information: Fentanyl Oral Transmucosal. www.Medlineplus.gov. January 01, 2006.
10. National Comprehensive Cancer Network. Practice Guidelines in Oncology; Adult Cancer Pain. 2.2005
11. Simmonds Mary. Management of Breakthrough Pain Due to Cancer. *Oncology*. August 1999. 13(8).
12. USP DI United States Pharmacopeia Dispensing Information (2006) volume 1. Thomson-Micromedex.
13. Actiq: <http://www.accessdata.fda.gov/scripts/cder/drugsatfda/index.cmf>
14. Fentora: <http://www.accessdata.fda.gov/scripts/cder/drugsatfda/index.cmf>
15. Cephalon, Inc. Actiq® Product Information. Cephalon, Inc.: Frazer, PA. February 2007.
16. Farrar JT, Cleary J, Rauck R: Oral transmucosal fentanyl citrate: randomized, double-blinded, placebo-controlled trial for treatment of breakthrough pain in cancer patients, *J Natl Cancer Inst* 1998 90: 611-6.
17. Landy SH. Oral transmucosal fentanyl citrate for the treatment of migraine headache pain in outpatients: a case series. *Headache*. 2004;44:762-6.
18. McEvoy GK, editor. AHFS Drug Information (online). American Society of Health-Systems Pharmacists, Bethesda, Maryland; 2006.
19. Oral transmucosal fentanyl citrate used to reduce emergency department visits in migraine patients: a prospective open label trial. [abstract]. *Pain Medicine* 2003;4:104. Abstract no. 536.
20. Shaiova L, Lapin J, Manco LS, et al. Tolerability and effects of two formulations of oral transmucosal fentanyl citrate (OTFC; ACTIQ) in patients with radiation-induced oral mucositis. *Support Care Cancer*. 2004 April;12(4):268-73.
21. Stein J. The American Academy of Pain Medicine (AAPM): Oral transmucosal fentanyl citrate helps migraineurs avoid emergency room. February 21, 2003 (cited January 5th, 2006). Available at: <http://www.docguide.com/news/content.nsf/news/8525697700573E1885256CD40059AF3F?OpenDocument&id=48DDE4A73E09A969852568880078C249&c=Pain&count=10>.

Coding

The five character codes included in the Blue Cross Blue Shield of Louisiana Medical Policy Coverage Guidelines are obtained from Current Procedural Terminology (CPT®)†, copyright 2013 by the American Medical Association (AMA). CPT is developed by the AMA as a listing of descriptive terms and five character identifying codes and modifiers for reporting medical services and procedures performed by physician.

The responsibility for the content of Blue Cross Blue Shield of Louisiana Medical Policy Coverage Guidelines is with Blue Cross and Blue Shield of Louisiana and no endorsement by the AMA is intended or should be implied. The AMA disclaims responsibility for any consequences or liability attributable or related to any use, nonuse or interpretation of information contained in Blue Cross Blue Shield of Louisiana Medical Policy Coverage Guidelines. Fee schedules, relative value units, conversion factors and/or related components are not assigned by the AMA, are not part of CPT,

©2014 Blue Cross and Blue Shield of Louisiana

An independent licensee of the Blue Cross and Blue Shield Association

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



BlueCross BlueShield of Louisiana

An independent licensee of the Blue Cross and Blue Shield Association.

Fentanyl Oral Transmucosal and Nasal Opioid Analgesics

Policy # 00213
Original Effective Date: 12/20/2006
Current Effective Date: 06/18/2014

and the AMA is not recommending their use. The AMA does not directly or indirectly practice medicine or dispense medical services. The AMA assumes no liability for data contained or not contained herein. Any use of CPT outside of Blue Cross Blue Shield of Louisiana Medical Policy Coverage Guidelines should refer to the most current Current Procedural Terminology which contains the complete and most current listing of CPT codes and descriptive terms. Applicable FARS/DFARS apply.

CPT is a registered trademark of the American Medical Association.

Codes used to identify services associated with this policy may include (but may not be limited to) the following:

| Code Type | Code |
|-----------------|------------------------|
| CPT | No codes |
| HCPCS | J3490 |
| ICD-9 Diagnosis | All relative diagnoses |
| ICD-9 Procedure | No codes |

Policy History

Original Effective Date: 12/20/2006
Current Effective Date: 06/18/2014

12/13/2006 Medical Director review
12/20/2006 Medical Policy Committee approval
12/12/2007 Medical Director review
12/19/2007 Medical Policy Committee approval. No change to coverage eligibility.
12/03/2008 Medical Director review
12/17/2008 Medical Policy Committee approval. Added note to coverage section from FDA regarding age restrictions for use of Actiq and Fentora.

11/12/2009 Medical Director review
11/18/2009 Medical Policy Committee approval. No change to coverage eligibility. Onsolis added to policy.
11/04/2010 Medical Policy Committee review
11/16/2010 Medical Policy Implementation Committee approval. No change to coverage eligibility.
03/01/2011 Added a note for the use of Abstral in the coverage section.
11/03/2011 Medical Policy Committee review
11/16/2011 Medical Policy Implementation Committee approval. Coverage eligibility unchanged.
02/02/2012 Medical Policy Committee review
02/15/2012 Medical Policy Implementation Committee approval. New nasal spray added to policy.
06/14/2012 Medical Policy Committee review
06/20/2012 Medical Policy Implementation Committee approval. Added the drug Subsys for use with age restrictions to the *Note* following the criteria in the coverage section.

03/04/2013 Coding update
06/06/2013 Medical Policy Committee review
06/25/2013 Medical Policy Implementation Committee approval. No change to coverage.
06/05/2014 Medical Policy Committee review
06/18/2014 Medical Policy Implementation Committee approval. Coverage eligibility unchanged.
Next Scheduled Review Date: 06/2015



BlueCross BlueShield of Louisiana

An independent licensee of the Blue Cross and Blue Shield Association.

Fentanyl Oral Transmucosal and Nasal Opioid Analgesics

Policy # 00213

Original Effective Date: 12/20/2006

Current Effective Date: 06/18/2014

***Investigational** – A medical treatment, procedure, drug, device, or biological product is Investigational if the effectiveness has not been clearly tested and it has not been incorporated into standard medical practice. Any determination we make that a medical treatment, procedure, drug, device, or biological product is Investigational will be based on a consideration of the following:

- A. whether the medical treatment, procedure, drug, device, or biological product can be lawfully marketed without approval of the U.S. Food and Drug Administration (FDA) and whether such approval has been granted at the time the medical treatment, procedure, drug, device, or biological product is sought to be furnished; or
- B. whether the medical treatment, procedure, drug, device, or biological product requires further studies or clinical trials to determine its maximum tolerated dose, toxicity, safety, effectiveness, or effectiveness as compared with the standard means of treatment or diagnosis, must improve health outcomes, according to the consensus of opinion among experts as shown by reliable evidence, including:
 1. Consultation with the Blue Cross and Blue Shield Association technology assessment program (TEC) or other nonaffiliated technology evaluation center(s);
 2. credible scientific evidence published in peer-reviewed medical literature generally recognized by the relevant medical community; or
 3. reference to federal regulations.

****Medically Necessary (or "Medical Necessity")** - Health care services, treatment, procedures, equipment, drugs, devices, items or supplies that a Provider, exercising prudent clinical judgment, would provide to a patient for the purpose of preventing, evaluating, diagnosing or treating an illness, injury, disease or its symptoms, and that are:

- A. in accordance with nationally accepted standards of medical practice;
- B. clinically appropriate, in terms of type, frequency, extent, level of care, site and duration, and considered effective for the patient's illness, injury or disease; and
- C. not primarily for the personal comfort or convenience of the patient, physician or other health care provider, and not more costly than an alternative service or sequence of services at least as likely to produce equivalent therapeutic or diagnostic results as to the diagnosis or treatment of that patient's illness, injury or disease.

For these purposes, "nationally accepted standards of medical practice" means standards that are based on credible scientific evidence published in peer-reviewed medical literature generally recognized by the relevant medical community, Physician Specialty Society recommendations and the views of Physicians practicing in relevant clinical areas and any other relevant factors.

‡ Indicated trademarks are the registered trademarks of their respective owners.

NOTICE: Medical Policies are scientific based opinions, provided solely for coverage and informational purposes. Medical Policies should not be construed to suggest that the Company recommends, advocates, requires, encourages, or discourages any particular treatment, procedure, or service, or any particular course of treatment, procedure, or service.