



BlueCross BlueShield of Louisiana

An independent licensee of the Blue Cross and Blue Shield Association.

Diagnosis and Treatment of Chronic Cerebrospinal Venous Insufficiency in Multiple Sclerosis

Policy # 00312

Original Effective Date: 08/17/2011

Current Effective Date: 09/17/2014

Applies to all products administered or underwritten by Blue Cross and Blue Shield of Louisiana and its subsidiary, HMO Louisiana, Inc. (collectively referred to as the "Company"), unless otherwise provided in the applicable contract. Medical technology is constantly evolving, and we reserve the right to review and update Medical Policy periodically.

Services Are Considered Investigational

Coverage is not available for investigational medical treatments or procedures, drugs, devices or biological products.

Based on review of available data, the Company considers the identification and subsequent treatment of chronic cerebrospinal venous insufficiency (CCSVI) in patients with multiple sclerosis (MS) to be **investigational**.*

Background/Overview

Chronic cerebrospinal venous insufficiency may be associated with MS, although this is controversial and an active area of research. Correction of CCSVI has been attempted via percutaneous venoplasty. The intent of this procedure is to relieve MS symptoms by improving venous drainage of the central nervous system. Correction of CCSVI by this method may be referred to as the "Liberation Procedure."

Multiple sclerosis is generally considered a chronic inflammatory demyelinating disease of the central nervous system (brain, spinal cord, optic nerve) felt to be triggered by an autoimmune response to myelin. However, in part due to the periventricular predilection of the lesions of multiple sclerosis, vascular etiologies CCSVI have also been considered. An animal model for MS was developed by injecting obstructing agents into the venous sinuses. This etiology, and treatment approach, for MS had not been actively pursued for many years; recent reports by a European researcher have renewed interest in this topic.

The core foundation of this vascular theory is that there is abnormal venous drainage from the brain due to outflow obstruction in the draining jugular vein and/or azygos veins. This abnormal venous drainage, which is characterized by special ultrasound criteria, is said to cause intracerebral flow disturbance or outflow problems that lead to periventricular deposits. In the CCSVI theory, these deposits have a similarity to the iron deposits seen around the veins in the legs in patients with chronic deep vein thrombosis. Those studying this theory have promoted balloon dilatation, with or without stenting, to treat the outflow problems, thereby curing CCSVI and by the same token alleviating MS complaints.

The following 5 criteria were defined by Zamboni et al. as features of CCSVI. In order to make the diagnosis of CCSVI, at least 2 of the 5 criteria need to be present:

1. Reflux constantly present (for a duration > 0.8s) in the supine and upright positions at the level of an internal jugular or vertebral vein. This parameter was evaluated during a short breath-hold following normal breathing and not under Valsalva maneuver.
2. Reflux at the level of veins of the deep cerebral system (for a duration > 0.5s). This was evaluated with the patient in the sitting and supine positions, and venous flow was enhanced by inviting the patient to breath in.

©2014 Blue Cross and Blue Shield of Louisiana

An independent licensee of the Blue Cross and Blue Shield Association

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



BlueCross BlueShield of Louisiana

An independent licensee of the Blue Cross and Blue Shield Association.

Diagnosis and Treatment of Chronic Cerebrospinal Venous Insufficiency in Multiple Sclerosis

Policy # 00312

Original Effective Date: 08/17/2011

Current Effective Date: 09/17/2014

3. Stenosis (< 0.3cm), valve abnormalities and septa on B-mode imaging.
4. Absence of flow at the level of the internal jugular or vertebral vein despite numerous deep inspirations.
5. No increase in the diameter of the internal jugular vein when changing from an upright to a supine position (lack of Δ -).

Centers for Medicare and Medicaid Services (CMS)

No national coverage determination (NCD) was identified. In the absence of an NCD, coverage decisions are left to the discretion of local Medicare carriers.

Rationale/Source

Diagnosis of Chronic Cerebrospinal Venous Insufficiency and Association with Multiple Sclerosis

Interest in the role of CCSVI in MS followed reports from a European vascular surgeon, Zamboni. Zamboni et al. used ultrasound and catheter-based venography to describe venous insufficiency in the internal jugular veins, vertebral veins, and deep cerebral veins of MS patients and described the finding as CCSVI. Using 5 ultrasound criteria (previously described), Zamboni defined the condition of CCSVI. In initial research reports, Zamboni et al reported both sensitivity and specificity of 100% in separating MS patients from controls when applying these criteria.

Since the development of the Zamboni criteria for CCSVI, there have been numerous research studies that have attempted to compare the rate of CCSVI in MS, normal patients, and patients with other neurologic diagnoses. The following review includes some of the largest of these studies, as well as any relevant systematic reviews and meta-analyses.

In 2014 a meta-analysis on the association between CCSVI and MS was published by Tsivgoulis et al. Included in the meta-analysis were 19 studies with a total of 1250 MS patients and 899 healthy controls. CCSVI was associated with MS in the pooled analysis with an odds ratio (OR) of 8.35 (95% confidence interval [CI], 3.44 to 20.31; $p < 0.001$). Heterogeneity across studies was considerable with a reported I^2 of 80.1%. In additional sensitivity analyses, the OR associating CCSVI and MS decreased. In the most conservative sensitivity analysis, which excluded 8 studies (studies by Zamboni or associated groups and studies conducted in Italy or by advocates of endovascular procedures for CCSVI), MS was not associated with CCSVI with an OR of 1.35 (95% CI, 0.62 to 2.93; $p = 0.453$). Additionally, heterogeneity was not found with a reported I^2 of 0%.

Zwischenberger et al in 2013 reported on a meta-analysis of the association between MS and CCSVI diagnosed by ultrasound. Included in the meta-analysis were 13 studies with a total of 1141 MS patients and 738 healthy controls. Initial analysis demonstrated CCSVI was associated with MS with an OR of 2.57 ($p < 0.001$), but heterogeneity was significant with a reported I^2 of 82.7% ($p < 0.001$). In a subsequent analysis of 9 studies with 4 outliers (studies with disproportionately high ORs) removed, the OR decreased, but still associated CCSVI with MS with an OR of 1.885 ($p < 0.001$) and heterogeneity decreased with an I^2 of 18 ($p = 0.279$).



BlueCross BlueShield of Louisiana

An independent licensee of the Blue Cross and Blue Shield Association.

Diagnosis and Treatment of Chronic Cerebrospinal Venous Insufficiency in Multiple Sclerosis

Policy # 00312

Original Effective Date: 08/17/2011

Current Effective Date: 09/17/2014

A systematic review of the association between CCSVI and MS was published in 2011 by Laupacis et al. This review included 8 studies that used ultrasound to diagnose CCSVI by the Zamboni criteria and compared the rate of CCSVI in patients with MS to those without MS. These studies were mostly small, with the median number of patients with MS of 50. There was a large degree of heterogeneity across studies in the rate of CCSVI among MS patients. Two smaller studies reported a rate of 0% for CCSVI in a total of 20 and 56 patients with MS. In contrast, the original study by Zamboni et al reported a 100% rate of CCSVI in 109 patients with MS. A small study of 25 patients also reported a very high rate of CCSVI at 84% (21/25). There was no obvious reason identified for this large discrepancy in CCSVI rates; the authors hypothesized that the most likely reason was variability in ultrasound technique and interpretation. There was a significant association of CCSVI with MS in combined analysis, with an odds ratio of 13.5 (95% CI, 2.6 to 71.4). There was a large amount of heterogeneity in this measure as well, with a reported I^2 of 89%. Several sensitivity analyses were performed, with marked variability of the OR from a low of 3.7 to more than 58,000, depending on the analysis. However, in all cases the association of CCSVI with MS remained significant.

A systematic review published in 2011 that included a smaller number of studies (n=4) came to similar conclusions. The rate of CCSVI in MS patients ranged from 7% to 100%, and the rate in non-MS patients ranged from 2% to 36%. There was a significant association between CCSVI but with a high degree of heterogeneity ($I^2=96%$) and an OR for association that had extreme variability, from approximately 2 to more than 26,000.

The largest study performed to date is a U.S. study by Zivadinov et al that used ultrasound to evaluate CCSVI in 499 subjects. A subject was considered CCSVI-positive if 2 or more venous hemodynamic (VH) criteria were fulfilled. The authors' studies of transcranial and extracranial echo-colored Doppler were carried out in 499 enrolled subjects: 289 with MS, 163 healthy controls (HC), 26 other neurologic diseases (OND), and 21 with clinically isolated syndromes (CIS). Prevalence rates for CCSVI were calculated in 3 ways: first, using only the subjects for whom diagnosis was certain (ie, borderline subjects were excluded); second, including the borderline subjects in the no CCSVI group; and finally, taking into account subjects who presented any of the VH criteria. CCSVI prevalence with borderline cases included in the no CCSVI group was 56.1% in MS, 42.3% in OND, 38.1% in CIS, and 22.7% in HC ($p<0.001$). The CCSVI prevalence figures were 62.5% for MS, 45.8% for OND, 42.1% for CIS, and 25.5% for HC when borderline cases were excluded ($p<0.001$). The prevalence of 1 or more positive VH criteria was the highest in MS (81.3%), followed by CIS (76.2%), OND (65.4%), and HC (55.2%) ($p<0.001$). CCSVI prevalence was higher in patients with progressive than in nonprogressive MS ($p=0.004$). The authors concluded that their findings were consistent with an increased prevalence of CCSVI in MS but with modest sensitivity and specificity. They also noted that their findings point against CCSVI having a primary causative role in the development of MS.

Zivadinov et al also reported on a substudy from the original study to explore any relationship between CCSVI and intracranial MS pathology as determined by magnetic resonance imaging (MRI). This substudy included 228 MS patients (162 relapsing-remitting and 66 secondary-progressive MS subtypes) and 73 HCs who had MRI imaging within 30 days of ultrasound imaging for CCSVI. In the MS group, 131 (57.5%) patients were considered CCSVI-positive and 21 (9.2%) were considered having borderline CCSVI. In the HC group, 19 (26%) were CCSVI-positive and 6 (8.2%) had borderline CCSVI. CCSVI was not significantly

©2014 Blue Cross and Blue Shield of Louisiana

An independent licensee of the Blue Cross and Blue Shield Association

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



BlueCross BlueShield of Louisiana

An independent licensee of the Blue Cross and Blue Shield Association.

Diagnosis and Treatment of Chronic Cerebrospinal Venous Insufficiency in Multiple Sclerosis

Policy # 00312
Original Effective Date: 08/17/2011
Current Effective Date: 09/17/2014

correlated with MRI imaging results on lesion burden and brain atrophy in MS patients and HCs. There was also no association between CCSVI and MRI imaging markers of inflammatory and neurodegenerative processes.

In an additional report from the Zivadinov study, Weinstock et al analyzed data from the MS subjects to examine the association between CCSVI and disability status, as measured by the Kurtzke Expanded Disability Status Scale (EDSS) and MS Severity Scale. CCSVI was not associated with disability status. However, there was an association between CCSVI and secondary or progressive MS versus nonprogressive MS, which included relapsing-remitting MS ($p=0.004$, $OR=2.34$; CI , 1.3 to 4.2).

Barreto et al conducted a single-center, prospective, case-control study of 206 MS and 70 non-MS patients to examine rates of CCSVI using color and spectral Doppler, B-mode imaging, with neurosonologists blinded to patients' clinical characteristics. Rates of CCSVI and extracranial or intracranial venous flow rates were not significantly different between MS and non-MS patients. In MS patients, CCSVI was found in 3.88% versus 7.14% of non-MS patients.

Floris et al used the Zamboni criteria to assess 74 patients with a diagnosis of MS and 34 healthy controls. All patients underwent Doppler ultrasound of the neck and transcranial Doppler ultrasound. A total of 34 patients were identified with CCSVI. The rate of CCSVI in MS patients was numerically higher than in normal controls (55% vs 35%), but this difference did not reach statistical significance ($p=0.09$). There were 12/74 patients 16% in the MS group who had normal ultrasound exams and an additional 28% (21 patients) who had some abnormalities on ultrasound but did not meet the criteria for CCSVI.

Centonze et al evaluated CCSVI by the Zamboni criteria in 84 patients with MS and 56 healthy controls. The rate of CCSVI was 50% in the MS patients versus 36% in controls ($p=0.12$). These authors also reported that there were no differences between MS patients that did and did not meet the criteria for CCSVI on demographic and clinical characteristics. There were also no differences between MS patients that did and did not meet CCSVI criteria in terms of disease severity, functional status, or quality of life. Doepp et al evaluated 56 patients with MS and 20 controls and found that none met the criteria for CCSVI using ultrasound.

Section Summary

The relationship between CCSVI and MS is unclear. The initial reports of excellent discrimination of MS patients from non-MS patients using CCSVI ultrasound criteria have not been replicated in subsequent studies. There is an extremely large variability in the literature in the rate of CCSVI among MS patients, ranging from 0% to 100%. Many of these studies report higher rates of CCSVI in MS patients, but others do not. Systematic reviews and meta-analyses have reported that the combined OR for an association is significantly increased; however, there is a very large degree of heterogeneity in these studies that has not been explained. If there is an association, it is unclear whether this is a causative factor for MS or whether the ultrasound findings are a result of MS and/or related processes.



BlueCross BlueShield of Louisiana

An independent licensee of the Blue Cross and Blue Shield Association.

Diagnosis and Treatment of Chronic Cerebrospinal Venous Insufficiency in Multiple Sclerosis

Policy # 00312

Original Effective Date: 08/17/2011

Current Effective Date: 09/17/2014

Treatment of CCSVI with Percutaneous Venoplasty

There are no trials with concurrent controls that report on outcomes of percutaneous venoplasty compared with alternatives. There is one small controlled trial of 15 patients comparing immediate venoplasty versus delayed venoplasty, and there are several single-arm case series that report on outcomes and adverse events of this procedure. One review and some of the larger case series are reviewed next.

In a 2012 Cochrane review, Van Zuuren et al found no randomized controlled trials (RCTs) on the treatment of CCSVI in MS patients. While there are ongoing clinical trials, the reviewers concluded the efficacy or safety of percutaneous transluminal angioplasty (PTA) for CCSVI treatment in MS patients could not be determined.

Hubbard et al prospectively followed 259 MS patients treated with venous angioplasty for CCSVI. Patients completed the Multiple Sclerosis Impact Scale (MSIS-29) 1 month before, and 1 and 6 months after angioplasty. MSIS scores significantly improved at each evaluation after angioplasty on both the physical and psychologic scales ($p < 0.01$). Symptoms recurred in 15 patients (6.3%).

In a case series of 65 patients with MS and CCSVI, Zamboni et al reported clinical improvement following catheter-based venoplasty. Patients were subdivided by MS clinical course into relapsing remitting ($n=35$), secondary progressive ($n=20$), and primary progressive ($n=10$) MS, and all patients underwent PTA. Mean follow-up was 18 months. In this study, outpatient endovascular treatment of CCSVI was noted to be feasible, with a minor complication rate. Postoperative venous pressure was significantly lower. The endovascular treatment was noted to improve MS clinical outcome measures, especially in the relapsing remitting group: the rate of relapse-free patients changed from 27% to 50% postoperatively ($p < 0.001$). The Multiple Sclerosis Functional Composite at 1 year improved significantly in relapsing remitting patients ($p < 0.008$) but not in primary progressive or secondary progressive. Physical quality of life (QOL) improved significantly in relapsing remitting ($p < 0.01$) and in primary progressive patients ($p < 0.03$), with a positive trend in secondary progressive ($p < 0.08$). The authors concluded that PTA of venous strictures in patients with CCSVI is safe, and especially in patients with relapsing remitting disease, the clinical course was positively influenced by treatment. The authors also indicated these results were from a pilot study and that a subsequent RCT is warranted.

Zamboni et al also reported a smaller series of 8 patients with ultrasound criteria for CCSVI undergoing immediate venoplasty compared with 7 patients undergoing delayed venoplasty. There were improvements on the EDSS for both groups following treatment, but no difference between groups in the first 6 months comparing immediate versus delayed treatment subjects. The relapse rate during the initial 6 months was 0.12% in the treatment group versus 0.66% in the control group, but this difference did not meet statistical significance. There were also trends toward improvement for the immediate treatment group on MRI scans, such as the number of T2 lesions, but these differences also did not reach statistical significance. No short-term adverse events were reported following the procedure, but the rate of restenosis at 1 year was 27% in treated patients.



BlueCross BlueShield of Louisiana

An independent licensee of the Blue Cross and Blue Shield Association.

Diagnosis and Treatment of Chronic Cerebrospinal Venous Insufficiency in Multiple Sclerosis

Policy # 00312

Original Effective Date: 08/17/2011

Current Effective Date: 09/17/2014

Adverse events: The initial small case series of venoplasty reported few adverse events. However, a number of larger case series have now been published that report on complications following endovascular interventions for CCSVI.

Burton et al described 5 patients who had undergone venoplasty and presented with complications of the procedure. The complications were internal jugular vein stent thrombosis, cerebral sinovenous thrombosis, stent migration, cranial nerve injury, and injury associated with venous catheterization. There was not a denominator in these studies to determine the rate of these events.

Petrov et al reported on the safety profile of 495 venoplasty procedures performed in 461 patients with MS, including 98 stent implantations. There were no deaths, major bleeding events, or acute exacerbations of MS. The most common procedure-related complication was vein dissection, which occurred in 3.0% of cases. Other complications included cardiac arrhythmias (1.2%), groin hematoma (1.0%), vein rupture (0.4%), and acute stent thrombosis (1.6%). Mandato et al reported adverse events within 30 days of endovascular intervention for 240 patients with MS over an 8-month period. Neck pain occurred in 15.6% of patients, most commonly following stent implantation. Headache occurred in 8.2% of patients and was persistent past 30 days in 1 patient (0.4%). Intraprocedural arrhythmias occurred in 1.3%, and 1 patient was diagnosed with a stress-induced cardiomyopathy following the procedure.

A U.S. Food and Drug Administration (FDA) alert was issued in May 2012 concerning the potential for adverse events following endovascular interventions for MS. Reports of adverse events obtained by FDA included death, stroke, detachment and/or migration of stents, vein damage, thrombosis, cranial nerve damage, and abdominal bleeding. This alert included the caveat that clinical trials of this procedure require FDA approval and an investigational device exemption because of the potential for harms.

Section Summary

The efficacy of venoplasty for CCSVI has been evaluated only in small case series and 1 very small trial of immediate versus delayed treatment. These studies report improvements in symptoms and disease-specific QOL measures. However, this evidence is insufficient to determine the efficacy of venoplasty because of the small amount of literature and the lack of controlled studies. RCTs are needed to adequately assess efficacy, especially when subjective patient-reported outcomes are used as the primary end point.

A few case series of several hundred patients have reported on adverse events. These studies establish that adverse events are uncommon following venoplasty, but serious adverse events do occur. FDA issued an alert in May 2012, noting the existence of serious complications, including death, and the need for ongoing monitoring. It is not currently possible to estimate the rate of serious adverse events such as death or major bleeding with confidence.

Ongoing Clinical Trials

A review of online site, ClinicalTrials.gov on April 24, 2014 identified 1 active phase 3 study. The BRAVE-DREAMS (Brain Venous Drainage Exploited Against Multiple Sclerosis) study is a multicenter, randomized, parallel group, blinded, sham-controlled trial to compare percutaneous balloon angioplasty with sham



BlueCross BlueShield of Louisiana

An independent licensee of the Blue Cross and Blue Shield Association.

Diagnosis and Treatment of Chronic Cerebrospinal Venous Insufficiency in Multiple Sclerosis

Policy # 00312

Original Effective Date: 08/17/2011

Current Effective Date: 09/17/2014

catheter venography in 679 patients with relapsing-remitting MS or secondary progressive MS and CCSVI (NCT01371760). This study has an expected completion date of August 2014.

Summary

Chronic cerebrospinal venous insufficiency may be associated with MS, although this is controversial and an active area of research. Correction of CCSVI has been attempted via percutaneous venoplasty. The intent of this procedure is to relieve MS symptoms by improving venous drainage of the central nervous system. Correction of CCSVI by this method may be referred to as the "Liberation Procedure."

The association of CCSVI with MS is uncertain. The rate of CCSVI in MS patients varies widely in the literature for unclear reasons, from 0% to 100%. Some studies report higher rates of CCSVI in patients with MS compared with non-MS patients, but others do not. If there is an association between MS and CCSVI, it is not known whether this is a causative factor for MS or a secondary result of the disease. It also appears that CCSVI can occur in other disorders and is not specific for MS.

Treatment of CCSVI with endovascular interventions has been attempted, but controlled trials to determine efficacy are lacking. The currently available studies report improvement in patient-reported symptoms following treatment, but this evidence is not sufficient to establish efficacy. Adverse events occur at a low overall rate, but serious adverse events can occur, and FDA issued an alert in 2012 concerning the potential for serious adverse events with treatment of CCSVI.

References

1. Blue Cross and Blue Shield Association, Medical Policy Reference Manual, "Diagnosis and Treatment of Chronic Cerebrospinal Venous Insufficiency in Multiple Sclerosis", 8.01.56, 5:2014.
2. Zamboni P, Galeotti R, Menegatti E et al. Chronic cerebrospinal venous insufficiency in patients with multiple sclerosis. *J Neurol Neurosurg Psychiatry* 2009; 80(4):392-9.
3. Zamboni P, Galeotti R, Menegatti E et al. A prospective open-label study of endovascular treatment of chronic cerebrospinal venous insufficiency. *J Vasc Surg* 2009; 50(6):1348-58 e1-3.
4. Tsvigoulis G, Sergentanis TN, Chan A et al. Chronic cerebrospinal venous insufficiency and multiple sclerosis: a comprehensive meta-analysis of case-control studies. *Thromb Haemostasis* 2014; 114(2):114-36.
5. Zwischenberger BA, Beasley MM, Davenport DL et al. Meta-analysis of the correlation between chronic cerebrospinal venous insufficiency and multiple sclerosis. *Vasc Endovascular Surg* 2013; 47(8):620-4.
6. Laupacis A, Lillie E, Dueck A et al. Association between chronic cerebrospinal venous insufficiency and multiple sclerosis: a meta-analysis. *CMAJ* 2011; 183(16):E1203-12.
7. Thapar A, Lane T, Nicholas R et al. Systematic review of sonographic chronic cerebrospinal venous insufficiency findings in multiple sclerosis. *Phlebology* 2011; 26(8):319-25.
8. Zivadinov R, Marr K, Cutter G et al. Prevalence, sensitivity, and specificity of chronic cerebrospinal venous insufficiency in MS. *Neurology* 2011; 77(2):138-44.
9. Zivadinov R, Cutter G, Marr K et al. No association between conventional brain MR imaging and chronic cerebrospinal venous insufficiency in multiple sclerosis. *AJNR Am J Neuroradiol* 2012; 33(10):1913-7.
10. Weinstock-Guttman B, Ramanathan M, Marr K et al. Clinical correlates of chronic cerebrospinal venous insufficiency in multiple sclerosis. *BMC Neurol* 2012; 12:26.
11. Barreto AD, Brod SA, Bui TT et al. Chronic cerebrospinal venous insufficiency: Case-control neurosonography results. *Ann Neurol* 2012.
12. Floris R, Centonze D, Fabiano S et al. Prevalence study of chronic cerebrospinal venous insufficiency in patients with multiple sclerosis: preliminary data. *Radiol Med* 2012.
13. Centonze D, Floris R, Stefanini M et al. Proposed chronic cerebrospinal venous insufficiency criteria do not predict multiple sclerosis risk or severity. *Ann Neurol* 2011; 70(1):51-8.

©2014 Blue Cross and Blue Shield of Louisiana

An independent licensee of the Blue Cross and Blue Shield Association

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Blue Cross Blue Shield of Louisiana

An independent licensee of the Blue Cross and Blue Shield Association.

Diagnosis and Treatment of Chronic Cerebrospinal Venous Insufficiency in Multiple Sclerosis

Policy # 00312

Original Effective Date: 08/17/2011

Current Effective Date: 09/17/2014

14. Doepp F, Paul F, Valdueza JM et al. No cerebrocervical venous congestion in patients with multiple sclerosis. *Ann Neurol* 2010; 68(2):173-83.
15. van Zuuren EJ, Fedorowicz Z, Pucci E et al. Percutaneous transluminal angioplasty for treatment of chronic cerebrospinal venous insufficiency (CCSVI) in multiple sclerosis patients. *Cochrane Database Syst Rev* 2012; 12:CD009903.
16. Hubbard D, Ponec D, Gooding J et al. Clinical improvement after extracranial venoplasty in multiple sclerosis. *J Vasc Interv Radiol* 2012; 23(10):1302-8.
17. Zamboni P, Galeotti R, Weinstock-Guttman B et al. Venous angioplasty in patients with multiple sclerosis: results of a pilot study. *Eur J Vasc Endovasc Surg* 2012; 43(1):116-22.
18. Burton JM, Alikhani K, Goyal M et al. Complications in MS patients after CCSVI procedures abroad (Calgary, AB). *Can J Neurol Sci* 2011; 38(5):741-6.
19. Petrov I, Grozdinski L, Kaninski G et al. Safety profile of endovascular treatment for chronic cerebrospinal venous insufficiency in patients with multiple sclerosis. *J Endovasc Ther* 2011; 18(3):314-23.
20. Mandato KD, Hegener PF, Siskin GP et al. Safety of endovascular treatment of chronic cerebrospinal venous insufficiency: a report of 240 patients with multiple sclerosis. *J Vasc Interv Radiol* 2012; 23(1):55-9.
21. FDA News Release: FDA issues alert on potential dangers of unproven treatment for multiple sclerosis. Available online at: <http://www.fda.gov/NewsEvents/Newsroom/PressAnnouncements/ucm303538.htm?source=govdelivery>. Last accessed April, 2014.
22. Reekers JA, Lee MJ, Belli AM et al. Cardiovascular and Interventional Radiological Society of Europe commentary on the treatment of chronic cerebrospinal venous insufficiency. *Cardiovasc Intervent Radiol* 2011; 34(1):1-2.
23. Vedantham S, Benenati JF, Kundu S et al. Interventional endovascular management of chronic cerebrospinal venous insufficiency in patients with multiple sclerosis: a position statement by the Society of Interventional Radiology, endorsed by the Canadian Interventional Radiology Association. *J Vasc Interv Radiol* 2010; 21(9):1335-7.
24. NICE. Percutaneous venoplasty for chronic cerebrospinal venous insufficiency for multiple sclerosis. March 2012. Available online at: www.nice.org.uk/ipg420. Last accessed April, 2014.
25. Baracchini C, Valdueza JM, Del Sette M et al. CCSVI and MS: a statement from the European Society of neurosonology and cerebral hemodynamics. *J Neurol* 2012; 259(12):2585-9.

Coding

The five character codes included in the Blue Cross Blue Shield of Louisiana Medical Policy Coverage Guidelines are obtained from Current Procedural Terminology (CPT)^{®†}, copyright 2013 by the American Medical Association (AMA). CPT is developed by the AMA as a listing of descriptive terms and five character identifying codes and modifiers for reporting medical services and procedures performed by physician.

The responsibility for the content of Blue Cross Blue Shield of Louisiana Medical Policy Coverage Guidelines is with Blue Cross and Blue Shield of Louisiana and no endorsement by the AMA is intended or should be implied. The AMA disclaims responsibility for any consequences or liability attributable or related to any use, nonuse or interpretation of information contained in Blue Cross Blue Shield of Louisiana Medical Policy Coverage Guidelines. Fee schedules, relative value units, conversion factors and/or related components are not assigned by the AMA, are not part of CPT, and the AMA is not recommending their use. The AMA does not directly or indirectly practice medicine or dispense medical services. The AMA assumes no liability for data contained or not contained herein. Any use of CPT outside of Blue Cross Blue Shield of Louisiana Medical Policy Coverage Guidelines should refer to the most current Current Procedural Terminology which contains the complete and most current listing of CPT codes and descriptive terms. Applicable FARS/DFARS apply.

CPT is a registered trademark of the American Medical Association.

Codes used to identify services associated with this policy may include (but may not be limited to) the following:

Code Type	Code
CPT	35460, 35476, 75978
HCPCS	No codes

©2014 Blue Cross and Blue Shield of Louisiana

An independent licensee of the Blue Cross and Blue Shield Association

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



BlueCross BlueShield of Louisiana

An independent licensee of the Blue Cross and Blue Shield Association.

Diagnosis and Treatment of Chronic Cerebrospinal Venous Insufficiency in Multiple Sclerosis

Policy # 00312
Original Effective Date: 08/17/2011
Current Effective Date: 09/17/2014

ICD-9 Diagnosis	340
ICD-9 Procedure	No codes

Policy History

Original Effective Date: 08/17/2011
Current Effective Date: 09/17/2014
08/04/2011 Medical Policy Committee review
08/17/2011 Medical Policy Implementation Committee approval. New policy.
08/02/2012 Medical Policy Committee review
08/15/2012 Medical Policy Implementation Committee approval. Coverage eligibility unchanged.
09/05/2013 Medical Policy Committee review
09/18/2013 Medical Policy Implementation Committee approval. Coverage eligibility unchanged.
09/04/2014 Medical Policy Committee review
09/17/2014 Medical Policy Implementation Committee approval. Coverage eligibility unchanged.
Next Scheduled Review Date: 09/2015

*Investigational – A medical treatment, procedure, drug, device, or biological product is Investigational if the effectiveness has not been clearly tested and it has not been incorporated into standard medical practice. Any determination we make that a medical treatment, procedure, drug, device, or biological product is Investigational will be based on a consideration of the following:

- A. Whether the medical treatment, procedure, drug, device, or biological product can be lawfully marketed without approval of the U.S. FDA and whether such approval has been granted at the time the medical treatment, procedure, drug, device, or biological product is sought to be furnished; or
- B. Whether the medical treatment, procedure, drug, device, or biological product requires further studies or clinical trials to determine its maximum tolerated dose, toxicity, safety, effectiveness, or effectiveness as compared with the standard means of treatment or diagnosis, must improve health outcomes, according to the consensus of opinion among experts as shown by reliable evidence, including:
 1. Consultation with the Blue Cross and Blue Shield Association technology assessment program (TEC) or other nonaffiliated technology evaluation center(s);
 2. Credible scientific evidence published in peer-reviewed medical literature generally recognized by the relevant medical community; or
 3. Reference to federal regulations.

‡ Indicated trademarks are the registered trademarks of their respective owners.

NOTICE: Medical Policies are scientific based opinions, provided solely for coverage and informational purposes. Medical Policies should not be construed to suggest that the Company recommends, advocates, requires, encourages, or discourages any particular treatment, procedure, or service, or any particular course of treatment, procedure, or service.