



# BlueCross BlueShield of Louisiana

An independent licensee of the Blue Cross and Blue Shield Association.

## Multianalyte Assays with Algorithmic Analyses for Predicting Risk of Type 2 Diabetes

**Policy #** 00418

**Original Effective Date:** 04/23/2014

**Current Effective Date:** 04/23/2014

*Applies to all products administered or underwritten by Blue Cross and Blue Shield of Louisiana and its subsidiary, HMO Louisiana, Inc. (collectively referred to as the "Company"), unless otherwise provided in the applicable contract. Medical technology is constantly evolving, and we reserve the right to review and update Medical Policy periodically.*

### **Services Are Considered Investigational**

*Coverage is not available for investigational medical treatments or procedures, drugs, devices or biological products.*

Based on review of available data, the Company considers the use of multianalyte panels with algorithmic analysis (MAAA) for the prediction of type 2 diabetes to be **investigational**.\*

### **Background/Overview**

The PreDx<sup>®</sup> Diabetes Risk Score (DRS) is a MAAA that is intended to predict the 5-year risk of type 2 diabetes. It is composed of 7 serum biomarkers that are combined via a proprietary algorithm. The proposed use is to identify patients at greater risk of developing type 2 diabetes and to potentially target preventive interventions at patients with the highest risk.

Type 2 diabetes mellitus is a highly prevalent disorder that is associated with an extremely high degree of morbidity and mortality. The true prevalence of type 2 diabetes in the U.S is uncertain due to a lack of population screening, but an estimated prevalence of 8.2% was reported in 2006. The incidence has been increasing rapidly over the last several decades, and current trends indicate that this increase will continue. Projections have estimated that the prevalence in the U.S. will reach 11.5% in 2011, 13.5% in 2021, and 14.5% in 2031.

Therefore, there is an urgent public health need to counter this trend. The potential to improve outcomes and reduce costs by preventing the onset of diabetes is vast. In order to accomplish this, accurate risk prediction methods may be helpful to identify populations with the highest risk of diabetes. Identification of patients at high risk could then be followed by preventive interventions targeted at high-risk individuals.

### **Predicting risk of type 2 diabetes.**

There are a variety of known factors that predict risk of type 2 diabetes. The most direct are measures of glucose metabolism, such as fasting glucose, oral glucose tolerance testing (OGTT), and hemoglobin A<sub>1c</sub> (HbA<sub>1c</sub>). For patients with impaired fasting glucose or impaired glucose tolerance, there is a high rate of progression to diabetes. Approximately 10% of these patients will progress to diabetes each year, and by 10 years more than 50% will have progressed to diabetes.

Other risk factors for diabetes include family history, ethnicity, lifestyle factors, dietary patterns, and numerous different laboratory parameters. A history of diabetes in the immediate family has long been recognized as one of the strongest predictors of diabetes. Regarding ethnicity, the risk of diabetes is increased 1.34 times for blacks, 1.86 times for Hispanics, and 2.26 times for Asians. A sedentary lifestyle, cigarette smoking, and dietary patterns that include sweetened foods and beverages have all been positively associated with the development of diabetes. In addition, there are numerous nonglucose

©2014 Blue Cross and Blue Shield of Louisiana

An independent licensee of the Blue Cross and Blue Shield Association

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



# BlueCross BlueShield of Louisiana

An independent licensee of the Blue Cross and Blue Shield Association.

## Multianalyte Assays with Algorithmic Analyses for Predicting Risk of Type 2 Diabetes

Policy # 00418  
Original Effective Date: 04/23/2014  
Current Effective Date: 04/23/2014

laboratory parameters that are associated with the risk of diabetes. These include inflammatory markers, lipid markers, measures of endothelial dysfunction, sex hormones, and many others.

Formal risk prediction instruments have combined clinical, laboratory, and genetic information to improve and refine upon the predictive ability of single factors. Many different formal risk prediction models have been developed. These models vary in the number and type of factors examined, and in the intended use of the instrument. For example, some prediction instruments consider the full range of clinical, biochemical, and genetic factors to derive the most accurate predictive model. Others, such as the Indian Risk Score and the Griffin Risk Score, use easily available clinical information without any laboratory markers to facilitate implementation as a widespread screening tool in areas of low resources.

In general, the available models have been shown to have good predictive ability, but most of them have not been externally validated. There is some evidence that directly compares the predictive accuracy of different measures, but there is insufficient comparative research to determine the optimal model. There is evidence that different models have different accuracy depending on the population tested. Also, relatively simple models have performed similarly to more complex models, and genetic information seems to add little over readily available clinical and metabolic parameters.

### Interventions to prevent type 2 diabetes.

A number of intervention trials have established that both lifestyle interventions and medications are effective in preventing the onset of type 2 diabetes in high-risk individuals. These trials have selected patients at high risk for diabetes, but have used single or several clinical factors, such as impaired glucose metabolism as selection factors, rather than formal risk prediction instruments. The largest reduction in diabetes incidence has been found for intensive lifestyle interventions that combine exercise and diet. There is a lesser effect for interventions with a single component and for interventions with medications.

A Cochrane review on the efficacy of lifestyle interventions to prevent type 2 diabetes was published in 2008. This review included 8 randomized trials that compared exercise and dietary interventions to standard therapy in patients at high risk for diabetes. There was a 37% reduction in the incidence of diabetes for the intervention cohort when a combined diet/exercise intervention was used, but there were not significant effects noted for an exercise-only or a diet-only intervention.

Another systematic review and meta-analysis evaluated the efficacy of medications for preventing progression to type 2 diabetes. This review included 10 studies of oral hypoglycemic agents and 15 studies of injectable agents. Oral hypoglycemic agents and orlistat were found to be effective in reducing progression to diabetes compared with usual care. In the largest trials with follow-up of greater than 2 years, metformin (relative risk [RR], 0.69; 95% confidence interval [CI], 0.57 to 0.83), acarbose (RR=0.75; 95% CI, 0.63 to 0.90), troglitazone (RR=0.45; 95% CI, 0.25 to 0.83), and orlistat (hazard ratio [HR], 0.63; 95% CI, 0.46 to 0.86) were efficacious in decreasing diabetes incidence compared with placebo. Evidence for other medication such as statins, fibrates, antihypertensive agents, and estrogen was inconclusive.

The largest randomized trial of preventive interventions was the Diabetes Prevention Program trial. This trial enrolled 3234 obese patients with a high risk of diabetes as defined by body mass index (BMI) level, fasting



# BlueCross BlueShield of Louisiana

An independent licensee of the Blue Cross and Blue Shield Association.

Multianalyte Assays with Algorithmic Analyses for Predicting Risk of Type 2 Diabetes

Policy # 00418  
Original Effective Date: 04/23/2014  
Current Effective Date: 04/23/2014

glucose, and 2-hour postprandial glucose levels. Participants were randomized to 1 of 3 groups, an intensive lifestyle intervention, a medication intervention consisting of metformin (850 mg twice per day), or a placebo control with information provided on diet and exercise. After a mean follow-up of 3 years, the incidence of diabetes was significantly reduced by 58% in the intensive lifestyle intervention group, and by 31% in the metformin group. A follow-up observational study concluded that the bulk of the benefit persisted for at least 10 years following completion of the trial.

## **PreDx Diabetes Risk Score**

The PreDx Diabetes Risk Score™<sup>‡</sup> (Tethys Bioscience®<sup>‡</sup> Inc., Emeryville, CA) is a commercially available MAAA that is intended to determine the 5-year risk of developing type 2 diabetes. The risk score is based on 7 biomarkers that are obtained by a peripheral blood draw:

- HgA<sub>1c</sub>
- Glucose
- Insulin
- C-reactive protein
- Ferritin
- Adiponectin
- Interleukin-2 receptor alpha

The results of these biomarkers are combined with age and gender to produce a quantitative risk score that varies from 0 to 10. Results are reported as the absolute 5-year risk of developing type 2 diabetes and the relative risk compared with age and gender matched controls.

## **FDA or Other Governmental Regulatory Approval**

U.S. Food and Drug Administration (FDA)

The biomarkers included in the PreDx Diabetes Risk Score are not subject to U.S. FDA approval. Laboratories performing these tests are subject to Clinical Laboratory Improvement Amendment standards for laboratory testing.

Centers for Medicare and Medicaid Services (CMS)

There is no national coverage determination (NCD). In the absence of an NCD, coverage decisions are left to the discretion of local Medicare carriers.

## **Rationale/Source**

### **Prediction of type 2 diabetes**

*Does the PreDx Diabetes Risk Score improve the ability to predict progression to diabetes, compared with standard clinical measurements?*

The development and validation of the PreDx DRS has been described in a series of manufacturer-sponsored studies. Kolberg et al first described the derivation of this risk score in 2009, using the Danish Inter99 patient cohort. This cohort consists of 61,031 subjects aged 30 to 60 years old and was intended to estimate the 5-year risk of progression to type 2 diabetes. The authors identified 64 candidate biomarkers that had support in the literature and that met study quality control indicators. They applied multiple logistic

©2014 Blue Cross and Blue Shield of Louisiana

An independent licensee of the Blue Cross and Blue Shield Association

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



# BlueCross BlueShield of Louisiana

An independent licensee of the Blue Cross and Blue Shield Association.

## Multianalyte Assays with Algorithmic Analyses for Predicting Risk of Type 2 Diabetes

Policy # 00418

Original Effective Date: 04/23/2014

Current Effective Date: 04/23/2014

regression approaches to select the biomarkers with the greatest predictive ability. Validation of the model with the same cohort was performed by the bootstrapping method.

The final model included 6 biomarkers: glucose, insulin, C-reactive protein, ferritin, adiponectin, and Interleukin-2 receptor alpha. The area under the curve (AUC) for the final fitted model was 0.78, and the bootstrapping estimate for AUC was similar at 0.76. The risk score was compared with single variables and simple combinations of variables. The best single predictor was the oral glucose tolerance test with an AUC of 0.79, which was not significantly different from the diabetes risk score. For the other single or combined variables, the AUC varied from 0.65 to 0.75. The diabetes risk score was superior to other single or combination variables, except for the 2-hour insulin level, which was not significantly different.

In a separate publication by the same research group using the same overall population of the Inter99 cohort, the model was validated in a different way. In this nested case-control design, 202 participants who progressed to type 2 diabetes were compared with 597 controls randomly selected from all participants who did not progress. The PreDx logistic model in this study consisted of the previously derived 6 biomarkers with the addition of HgA<sub>1c</sub>.

The AUC of the fitted model was 0.84. This was superior to single biomarkers, which had AUCs that ranged from 0.70 to 0.77, and was also superior to a noninvasive clinical model that had an AUC of 0.77. The absolute 5-year risk of progression to diabetes for patients with a low DRS (<4.5) was 1%, which rose to 7% for patients with a moderate DRS (≥4.5 and <8), and to 24% for patients with a high DRS (≥8.0).

A second validation study used a separate cohort from the prospective Botnia study. This was a cohort of 2770 individuals who were at increased risk of developing type 2 diabetes due mainly to family history. Outcome data and biomarker data were available for 2350 individuals.

The AUC for the validation set was 0.78, which was lower than the AUC of 0.84 obtained for the training set. The absolute 5-year risk of progression to diabetes for patients with a low DRS (<4.5) was 1.1% (95% CI, 0.5% to 1.6%), which rose to 4.0% (95% CI, 2.3% to 5.7%) for patients with a moderate DRS (≥4.5 and <8), and to 12.7% (95% CI, 7.0% to 18.1%) for patients with a high DRS (≥8.0). Reclassification analysis was also performed using fasting glucose and OGTT as baseline. The net reclassification index of 0.20 indicated that the DRS performed better than glucose and OGTT. The main advantage of the DRS was in reclassifying patients with abnormal glucose and OGTT into lower risk levels after application of the DRS.

*How does the PreDx Diabetes Risk Score compare with other diabetes risk scores in predicting the future risk of type 2 diabetes?*

There is a body of literature on the comparative accuracy of different DRSs. However, the most studies that directly compare different risk scores do not include the PreDx DRS as one of the comparators. For example, Abbasi et al performed a systematic review and independent validation of 12 different risk models identified in the literature, but did not include the PreDx score. In another publication evaluating the validity of different risk models a total of 5 risk scores were reviewed, but the PreDx score was not included. Noble et al conducted a systematic review of DRSs and included the Inter99 Danish cohort study score from



# BlueCross BlueShield of Louisiana

An independent licensee of the Blue Cross and Blue Shield Association.

## Multianalyte Assays with Algorithmic Analyses for Predicting Risk of Type 2 Diabetes

Policy # 00418  
Original Effective Date: 04/23/2014  
Current Effective Date: 04/23/2014

which the PreDx score was derived, but the authors do not specify that the PreDx score is specifically used. The authors make no direct comparisons between risk scores, noting that direct comparisons are precluded by heterogeneity in the patient populations, clinical outcomes reported, and intended context of use, among other factors.

Within the manufacturer-sponsored validation studies, there were limited comparisons of the PreDx score with other risk models. PreDx score was compared with single markers and simple combinations of markers and was superior to most of these comparators. Lyssenko et al compared the performance of the DRS with 2 other risk prediction scores, the Framingham score and the San Antonio Heart risk score. The Framingham score had an AUC of 0.76 while the San Antonio score had an AUC of 0.77, neither of which were significantly different from the DRS, which had an AUC of 0.78.

Subsequently, Rowe et al compared the PreDx score with other clinical variables used for diabetes risk prediction in a different patient population. The Insulin Resistance Atherosclerosis Study cohort was a multiethnic U.S. cohort, including 722 patients, which was designed to evaluate insulin resistance and cardiovascular risk factors and disease states in different U.S. ethnic groups and varying states of glucose tolerance. This study compared the 5-year risk of type 2 diabetes as estimated by the DRS with other risk assessment tools, including fasting glucose, BMI, fasting insulin, the homeostasis model assessment-estimated insulin resistance (HOMA-IR) index, and the OGTT. Test performance was assessed by receiver operator characteristic curve analysis, with AUC reported. In the whole cohort, the DRS had a significantly higher AUC than fasting glucose alone (0.762 vs 0.711,  $p=0.003$ ), fasting insulin alone (0.690,  $p=0.003$ ), HOMA-IR (0.716,  $p=0.03$ ), and BMI (0.671,  $p<0.001$ ). It was not statistically different from the 2-hour glucose tolerance test.

### Section summary

The PreDx DRS has been tested in 2 different prospective cohorts of patients, with reported AUCs of 0.78 and 0.84, indicating good overall accuracy for predicting progression to diabetes. It has been evaluated in 1 U.S. cohort, but has otherwise not been tested in a wide range of patient populations. As a result, there is some uncertainty in the predictive accuracy and generalizability of the risk score.

The evidence is insufficient to determine the comparative efficacy of the PreDx score compared with other diabetes risk scores. The single study that compared the PreDx score with 2 established measures (Framingham diabetes risk score, San Antonio Heart diabetes risk score) reported that the overall accuracy, as defined by AUC for predicting progression to diabetes, did not differ significantly among the 3 measures. A study in a U.S. cohort of patients suggested that the PreDx score may better predict diabetes than several individual risk factors alone. However, this comparative evidence is incomplete, and more comprehensive comparative studies are needed.

### Prevention of type 2 diabetes

*Does the use of MAAAs lead to targeted interventions that reduce the incidence of type 2 diabetes?*

In 2013, Shah et al published results from the industry-sponsored Provision of Evidence-based Therapies Among Individuals at High Risk for Type 2 Diabetes (PREVAIL) initiative, a retrospective cohort study



# BlueCross BlueShield of Louisiana

An independent licensee of the Blue Cross and Blue Shield Association.

## Multianalyte Assays with Algorithmic Analyses for Predicting Risk of Type 2 Diabetes

Policy # 00418  
Original Effective Date: 04/23/2014  
Current Effective Date: 04/23/2014

designed to evaluate the influence of the PreDx score on the use of interventions related to prediabetes and diabetes risk factors. The study included 30 sites across the U.S., each of which retrospectively abstracted chart information for up to 50 consecutive patients who had undergone PreDx testing at a baseline visit and a follow-up visit, for a total of 913 patients. The PreDx test score was stratified into low-, moderate-, and high-risk groups. From baseline to follow-up, all patients demonstrated small reductions in systolic blood pressure (from 128 to 165.5,  $p=0.039$ ), increased antihypertensive use among those with hypertension (from 73.1% to 77.2%,  $p<0.001$ ), decreased median low-density lipoprotein (LDL) (from 104 mg/dL to 100 mg/dL,  $p=0.009$ ), and increased median high-density lipoprotein (HDL) (from 48 mg/dL to 50 mg/dL,  $p<0.001$ ). A similar proportion of patients received lifestyle counseling at follow-up as at baseline. The PreDx risk group was not significantly associated with changes in systolic blood pressure, antihypertensive use, or changes in LDL or HDL. However, patients with higher PreDx risk groups were more likely to undergo lifestyle counseling. Limitations of this study include a lack of standardized inclusion criteria, lack of a comparison group, and nonstandardized use of the PreDx test for clinical decision making. Any interventions were left up to physician and/or patient discretion. These limitations make it impossible to determine what changes would have occurred in the absence of any intervention or with only routine medical care. Therefore, this study provides little evidence to support using the PreDx score as a means to select patients for diabetes-related interventions.

No other studies were identified that used the PreDx DRS as a method to select patients for interventions to prevent type 2 diabetes.

Other risk prediction instruments have been used for this purpose, demonstrating the potential for the use of risk prediction instruments to target preventive interventions. The AusDiab study used the Australian Type 2 Diabetes Risk Assessment Tool, based on 9 self-assessed measures, to evaluate the efficiency of detecting individuals at high risk of diabetes. This study used data from 5814 participants in the Australian Diabetes, Obesity, and Lifestyle Study to model 4 different screening strategies. The optimal strategy, defined as resulting in the greatest number of patients entered into preventive interventions at the least total costs, was noninvasive screening with the Diabetes Risk Assessment Tool followed by measurement of fasting plasma glucose.

### Section summary

The evidence is insufficient to determine whether the PreDx risk score can improve outcomes by targeting preventive interventions to patients who will benefit most. One study evaluated changes in cardiovascular risk factors in patients whose physicians used the PreDx score, but there are no published studies that evaluate use of the risk score to target preventive interventions. It is not known whether the PreDx risk score is as good as or better than other methods for identifying individuals at high risk for diabetes.

### Summary

The PreDx DRS is a multianalyte assay with algorithmic analysis that uses 7 biomarkers, including glucose and HgA<sub>1C</sub>, to predict progression to diabetes. There is limited published evidence on the validity and applicability of this risk score. Two patient cohorts reported area under the curve for predicting progression to diabetes that ranged from 0.78 to 0.84. This suggests good overall predictive ability, but is limited by the lack of validation by independent research groups and testing in a wider variety of patient populations. The



# BlueCross BlueShield of Louisiana

An independent licensee of the Blue Cross and Blue Shield Association.

## Multianalyte Assays with Algorithmic Analyses for Predicting Risk of Type 2 Diabetes

Policy # 00418

Original Effective Date: 04/23/2014

Current Effective Date: 04/23/2014

evidence is insufficient to determine the comparative accuracy of the PreDx DRS with other formal prediction models for diabetes.

There is a lack of evidence on the clinical utility of the PreDx score. No published studies were identified that used the risk score to select patients for preventive interventions. As a result, it is not known how this instrument will perform in targeting preventive interventions to patients who will benefit the most, nor is it known how this risk score compares with other methods for selecting high-risk patients. Therefore, use of the PreDx diabetes risk score is considered investigational.

### References

1. Blue Cross and Blue Shield Association, Medical Policy Reference Manual, "Multianalyte Assays with Algorithmic Analyses for Predicting Risk of Type 2 Diabetes", 2.04.90, 2:2014.
2. Fox CS, Pencina MJ, Meigs JB et al. Trends in the incidence of type 2 diabetes mellitus from the 1970s to the 1990s: the Framingham Heart Study. *Circulation* 2006; 113(25):2914-8.
3. Blonde L. State of diabetes care in the United States. *Am J Manag Care* 2007; 13 Suppl 2:S36-40.
4. Mainous AG, 3rd, Baker R, Koopman RJ et al. Impact of the population at risk of diabetes on projections of diabetes burden in the United States: an epidemic on the way. *Diabetologia* 2007; 50(5):934-40.
5. Schwarz PE, Li J, Lindstrom J et al. Tools for predicting the risk of type 2 diabetes in daily practice. *Horm Metab Res* 2009; 41(2):86-97.
6. Shai I, Jiang R, Manson JE et al. Ethnicity, obesity, and risk of type 2 diabetes in women: a 20-year follow-up study. *Diabetes Care* 2006; 29(7):1585-90.
7. Liu S, Tinker L, Song Y et al. A prospective study of inflammatory cytokines and diabetes mellitus in a multiethnic cohort of postmenopausal women. *Arch Intern Med* 2007; 167(15):1676-85.
8. Meigs JB, Hu FB, Rifai N et al. Biomarkers of endothelial dysfunction and risk of type 2 diabetes mellitus. *JAMA* 2004; 291(16):1978-86.
9. Noble D, Mathur R, Dent T et al. Risk models and scores for type 2 diabetes: systematic review. *BMJ* 2011; 343.
10. Balkau B, Lange C, Fezeu L et al. Predicting diabetes: clinical, biological, and genetic approaches: data from the Epidemiological Study on the Insulin Resistance Syndrome (DESIR). *Diabetes Care* 2008; 31(10):2056-61.
11. Mohan V, Goldhaber-Fiebert JD, Radha V et al. Screening with OGTT alone or in combination with the Indian diabetes risk score or genotyping of TCF7L2 to detect undiagnosed type 2 diabetes in Asian Indians. *Indian J Med Res* 2011; 133:294-9.
12. Griffin SJ, Little PS, Hales CN et al. Diabetes risk score: towards earlier detection of type 2 diabetes in general practice. *Diabetes Metab Res Rev* 2000; 16(3):164-71.
13. Abbasi A, Peelen LM, Corpeleijn E et al. Prediction models for risk of developing type 2 diabetes: systematic literature search and independent external validation study. *BMJ* 2012; 345:e5900.
14. Orozco LJ, Buchleitner AM, Gimenez-Perez G et al. Exercise or exercise and diet for preventing type 2 diabetes mellitus. *Cochrane Database Syst Rev* 2008; (3):CD003054.
15. Padwal R, Majumdar SR, Johnson JA et al. A systematic review of drug therapy to delay or prevent type 2 diabetes. *Diabetes Care* 2005; 28(3):736-44.
16. Knowler WC, Barrett-Connor E, Fowler SE et al. Reduction in the incidence of type 2 diabetes with lifestyle intervention or metformin. *N Engl J Med* 2002; 346(6):393-403.
17. Diabetes Prevention Program Research G, Knowler WC, Fowler SE et al. 10-year follow-up of diabetes incidence and weight loss in the Diabetes Prevention Program Outcomes Study. *Lancet* 2009; 374(9702):1677-86.
18. Kolberg JA, Jorgensen T, Gerwien RW et al. Development of a type 2 diabetes risk model from a panel of serum biomarkers from the Inter99 cohort. *Diabetes Care* 2009; 32(7):1207-12.
19. Lyssenko V, Jorgensen T, Gerwien RW et al. Validation of a multi-marker model for the prediction of incident type 2 diabetes mellitus: combined results of the Inter99 and Botnia studies. *Diab Vasc Dis Res* 2012; 9(1):59-67.
20. Urdea M, Kolberg J, Wilber J et al. Validation of a multimarker model for assessing risk of type 2 diabetes from a five-year prospective study of 6784 Danish people (Inter99). *J Diabetes Sci Technol* 2009; 3(4):748-55.
21. Rowe MW, Bergman RN, Wagenknecht LE et al. Performance of a multi-marker diabetes risk score in the Insulin Resistance Atherosclerosis Study (IRAS), a multi-ethnic US cohort. *Diabetes Metab Res Rev* 2012; 28(6):519-26.
22. Shah BR, Cox M, Inzucchi SE et al. A quantitative measure of diabetes risk in community practice impacts clinical decisions: The PREVAIL initiative. *Nutr Metab Cardiovasc Dis* 2013.

©2014 Blue Cross and Blue Shield of Louisiana

An independent licensee of the Blue Cross and Blue Shield Association

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



# Blue Cross Blue Shield of Louisiana

An independent licensee of the Blue Cross and Blue Shield Association.

## Multianalyte Assays with Algorithmic Analyses for Predicting Risk of Type 2 Diabetes

Policy # 00418  
 Original Effective Date: 04/23/2014  
 Current Effective Date: 04/23/2014

23. Chen L, Magliano DJ, Balkau B et al. Maximizing efficiency and cost-effectiveness of Type 2 diabetes screening: the AusDiab study. *Diabet Med* 2011; 28(4):414-23.
24. Association AD. Standards of Medical Care in Diabetes—2014. *Diabetes Care* 2014; 37(Supplement 1):S14-S80.
25. U. S. Preventive Services Task Force. Screening for type 2 diabetes mellitus in adults: U.S. Preventive Services Task Force recommendation statement. *Ann Intern Med* 2008; 148(11):846-54.

### Coding

The five character codes included in the Blue Cross Blue Shield of Louisiana Medical Policy Coverage Guidelines are obtained from Current Procedural Terminology (CPT®)†, copyright 2013 by the American Medical Association (AMA). CPT is developed by the AMA as a listing of descriptive terms and five character identifying codes and modifiers for reporting medical services and procedures performed by physician.

The responsibility for the content of Blue Cross Blue Shield of Louisiana Medical Policy Coverage Guidelines is with Blue Cross and Blue Shield of Louisiana and no endorsement by the AMA is intended or should be implied. The AMA disclaims responsibility for any consequences or liability attributable or related to any use, nonuse or interpretation of information contained in Blue Cross Blue Shield of Louisiana Medical Policy Coverage Guidelines. Fee schedules, relative value units, conversion factors and/or related components are not assigned by the AMA, are not part of CPT, and the AMA is not recommending their use. The AMA does not directly or indirectly practice medicine or dispense medical services. The AMA assumes no liability for data contained or not contained herein. Any use of CPT outside of Blue Cross Blue Shield of Louisiana Medical Policy Coverage Guidelines should refer to the most current Current Procedural Terminology which contains the complete and most current listing of CPT codes and descriptive terms. Applicable FARS/DFARS apply.

CPT is a registered trademark of the American Medical Association.

Codes used to identify services associated with this policy may include (but may not be limited to) the following:

Code Type	Code
CPT	81506
HCPCS	No codes
ICD-9 Diagnosis	All relative diagnoses
ICD-9 Procedure	No codes

### Policy History

Original Effective Date: 04/23/2014  
 Current Effective Date: 04/23/2014  
 04/03/2014 Medical Policy Committee review  
 04/23/2014 Medical Policy Implementation Committee approval. New policy.  
 Next Scheduled Review Date: 04/2015

\*Investigational – A medical treatment, procedure, drug, device, or biological product is Investigational if the effectiveness has not been clearly tested and it has not been incorporated into standard medical practice. Any determination we make that a medical treatment, procedure, drug, device, or biological product is Investigational will be based on a consideration of the following:

- A. whether the medical treatment, procedure, drug, device, or biological product can be lawfully marketed without approval of the U.S. FDA and whether such approval has been granted at the time the medical treatment, procedure, drug, device, or biological product is sought to be furnished; or
- B. whether the medical treatment, procedure, drug, device, or biological product requires further studies or clinical trials to determine its maximum tolerated dose, toxicity, safety, effectiveness, or effectiveness as compared with the standard means

©2014 Blue Cross and Blue Shield of Louisiana

An independent licensee of the Blue Cross and Blue Shield Association

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



# BlueCross BlueShield of Louisiana

An independent licensee of the Blue Cross and Blue Shield Association.

## Multianalyte Assays with Algorithmic Analyses for Predicting Risk of Type 2 Diabetes

Policy # 00418

Original Effective Date: 04/23/2014

Current Effective Date: 04/23/2014

of treatment or diagnosis, must improve health outcomes, according to the consensus of opinion among experts as shown by reliable evidence, including:

1. Consultation with the Blue Cross and Blue Shield Association technology assessment program (TEC) or other nonaffiliated technology evaluation center(s);
2. credible scientific evidence published in peer-reviewed medical literature generally recognized by the relevant medical community; or
3. reference to federal regulations.

‡ Indicated trademarks are the registered trademarks of their respective owners.

**NOTICE:** Medical Policies are scientific based opinions, provided solely for coverage and informational purposes. Medical Policies should not be construed to suggest that the Company recommends, advocates, requires, encourages, or discourages any particular treatment, procedure, or service, or any particular course of treatment, procedure, or service.