



## Deep Brain Stimulation

**Policy Number:** 7.01.63  
**Origination:** 8/2001

**Last Review:** 8/2014  
**Next Review:** 8/2015

### **Policy**

---

Blue Cross and Blue Shield of Kansas City (Blue KC) will provide coverage for deep brain stimulation when it is determined to be medically necessary because the criteria shown below are met.

### **When Policy Topic is covered**

---

Unilateral deep brain stimulation of the thalamus may be considered **medically necessary** in patients with disabling, medically unresponsive tremor due to essential tremor or Parkinson's disease.

Unilateral or bilateral deep brain stimulation of the globus pallidus or subthalamic nucleus may be considered **medically necessary** in the following patients:

- Those with Parkinson's Disease and ALL of the following:
  - a good response to levodopa; AND
  - a minimal score of 30 points on the motor portion of the Unified Parkinson Disease Rating Scale when the patient has been without medication for approximately 12 hours; AND
  - motor complications not controlled by pharmacologic therapy
- Patients aged greater than 7 years with chronic, intractable (drug refractory) primary dystonia, including generalized and/or segmental dystonia, hemidystonia and cervical dystonia (torticollis).

### **When Policy Topic is not covered**

---

Deep brain stimulation for other movement disorders, including but not limited to multiple sclerosis, post-traumatic dyskinesia, and tardive dyskinesia is considered **investigational**.

Deep brain stimulation for the treatment of chronic cluster headaches is considered **investigational**.

Deep brain stimulation for the treatment of other psychiatric or neurologic disorders, including but not limited to Tourette syndrome, depression, obsessive-compulsive disorder, Alzheimer disease, anorexia nervosa, alcohol addiction, chronic pain, and epilepsy, is considered **investigational**.

### **Contraindications** to deep brain stimulation include:

- Patients who are not good surgical risks because of unstable medical problems or because of the presence of a cardiac pacemaker
- Patients who have medical conditions that require repeated magnetic resonance imaging (MRI)
- Patients who have dementia that may interfere with the ability to cooperate
- Patients who have had botulinum toxin injections within the last 6 months

### **Considerations**

---

Disabling, medically unresponsive tremor is defined as tremor that causes significant limitation in daily activities and inadequate control by maximal dosage of medication for at least 3 months before implant. This includes problems related to the drugs causing serious side effects, or medications that cause no improvement or transient improvement.

## **Description of Procedure or Service**

---

Deep brain stimulation (DBS) involves the stereotactic placement of an electrode into the brain (i.e., hypothalamus, thalamus, globus pallidus, or subthalamic nucleus). DBS is used as an alternative to permanent neuroablative procedures for control of essential tremor (ET) and Parkinson's disease (PD). DBS is also being evaluated for the treatment of a variety of other neurologic and psychiatric disorders, including epilepsy, dystonia, cluster headache, Tourette syndrome, depression, and obsessive-compulsive disorder (OCD).

---

## **Background**

Deep brain stimulation (DBS) has been investigated as an alternative to permanent neuroablative procedures, such as thalamotomy and pallidotomy. The technique has been most thoroughly investigated as an alternative to thalamotomy for unilateral control of essential tremor (ET) and tremor associated with Parkinson's disease (PD). More recently, there has been research interest in the use of DBS of the globus pallidus or subthalamic nucleus as a treatment of other parkinsonian symptoms, such as rigidity, bradykinesia, or akinesia. Another common morbidity associated with PD is the occurrence of motor fluctuations, referred to as "on and off" phenomena, related to the maximum effectiveness of drugs (i.e., the "on" state) and the nadir response during drug troughs (i.e., the "off" state). In addition, levodopa, the most commonly used anti-Parkinson's drug, may be associated with disabling drug-induced dyskinesias. Therefore, the optimal pharmacologic treatment of PD may involve a balance between optimal effects on Parkinson's disease symptoms versus the appearance of drug-induced dyskinesias. The effect of DBS on both Parkinson's disease symptoms and drug-induced dyskinesias has also been studied.

DBS has also been investigated in patients with primary and secondary dystonia, defined as a neurologic movement disorder characterized by involuntary muscle contractions, which force certain parts of the body into abnormal, contorted, and painful movements or postures. Dystonia can be classified according to age of onset, bodily distribution of symptoms, and cause. Age of onset can occur during childhood or during adulthood. Dystonia can affect certain portions of the body (focal dystonia and multifocal dystonia) or the entire body (generalized dystonia). Torticollis is an example of a focal dystonia. Primary dystonia is defined when dystonia is the only symptom unassociated with other pathology. Treatment options for dystonia include oral or injectable medications (i.e., botulinum toxin) and destructive surgical or neurosurgical interventions (i.e., thalamotomies or pallidotomies) when conservative therapies fail. Secondary dystonia is a dystonia brought on by an inciting event, such as a stroke, trauma, or drugs. Tardive dystonia is a form of drug-induced secondary dystonia.

DBS has been investigated in patients with chronic cluster headaches. Cluster headaches occur as episodic attacks of severe pain lasting from 30 minutes to several hours. The pain is usually unilateral and localized to the eye, temple, forehead, and side of the face. Autonomic symptoms that occur with cluster headaches include ipsilateral facial sweating, flushing, tearing, and rhinorrhea. Cluster headaches occur primarily in men and have been classified as vascular headaches that have been associated with high blood pressure, smoking, alcohol use, etc. However, the exact pathogenesis of cluster headaches is uncertain. Positron emission tomography (PET) scanning and magnetic resonance imaging (MRI) have shown the hypothalamic region may be important in the pathogenesis of cluster headaches. Alterations in hormonal/serotonergic function may also play a role. Treatment of cluster headaches includes pharmacologic interventions for acute episodes and prophylaxis, sphenopalatine ganglion (SPG) blockade, and surgical procedures such as percutaneous SPG radiofrequency rhizotomy and gamma knife radiosurgery of the trigeminal nerve.

The role of DBS in treatment of other treatment-resistant neurologic and psychiatric disorders, particularly Tourette syndrome, epilepsy, obsessive-compulsive disorder (OCD), and major depressive disorders, is also being investigated. Ablative procedures are irreversible and, though they have been refined, remain controversial treatments for intractable illness. Interest has shifted to neuromodulation through DBS of nodes or targets within neural circuits involved in these disorders. Currently, a variety of target areas are being studied.

DBS involves the stereotactic placement of an electrode into the brain (i.e., hypothalamus, thalamus, globus pallidus, or subthalamic nucleus). The electrode is initially attached to a temporary transcutaneous cable for short-term stimulation to validate treatment effectiveness. Several days later, the patient returns to surgery for permanent subcutaneous implantation of the cable and a radiofrequency-coupled or battery-powered programmable stimulator. The electrode is typically implanted unilaterally on the side corresponding to the most severe symptoms. However, the use of bilateral stimulation using 2 electrode arrays has also been investigated in patients with bilateral, severe symptoms. After implantation, noninvasive programming of the neurostimulator can be adjusted to the patient's symptoms. This feature may be important for patients with PD, whose disease may progress over time, requiring different neurostimulation parameters. Setting the optimal neurostimulation parameters may involve the balance between optimal symptom control and appearance of side effects of neurostimulation, such as dysarthria, disequilibrium, or involuntary movements.

### **Regulatory Status**

The U.S. Food and Drug Administration (FDA) has approved the Activa® Tremor Control System, manufactured by Medtronic Corp, MN, for deep brain stimulation. While the original 1997 FDA-labeled indications were limited to unilateral implantation of the device for the treatment of tremor, in January 2002, the FDA-labeled indications were expanded to include bilateral implantation as a treatment to decrease the symptoms of advanced Parkinson's disease that are not controlled by medication. In April 2003, the labeled indications were expanded to include "unilateral or bilateral stimulation of the internal globus pallidus or subthalamic nucleus to aid in the management of chronic, intractable (drug refractory) primary dystonia, including generalized and/or segmental dystonia, hemidystonia, and cervical dystonia (torticollis) in patients seven years of age or above." This latter indication received FDA approval through the Humanitarian Device Exemption process. The Activa Tremor Control System consists of the following components: the implantable pulse generator, the deep brain stimulator lead, an extension that connects the lead to the power source, a console programmer, a software cartridge to set electrical parameters for stimulation, and a patient control magnet, which allows the patient to turn the pulse generator on and off, or change between high and low settings.

The Vercise™ Deep Brain Stimulation (DBS) system (Boston Scientific) is currently available in Europe, Israel and Australia. Completion of a large U.S. multicenter trial (INTREPID) is expected in 2021.

In February 2009, the FDA approved deep brain stimulation with the Reclaim® device (Medtronic, Inc.) via the Humanitarian Device Exemption (HDE) process for the treatment of severe obsessive-compulsive disorder (OCD).

### **Rationale**

This policy was updated with a literature review through June 11, 2014. The policy was originally based on 2 TEC Assessments; a 1997 TEC Assessment that focused on unilateral deep brain stimulation (DBS) of the thalamus as a treatment for tremor and a 2001 TEC Assessment that focused on the use of DBS of the globus pallidus and subthalamic nucleus for a broader range of Parkinson's disease (PD) symptoms. (1, 2) The observations and conclusions of the TEC Assessments are summarized here.

#### **Unilateral Deep Brain Stimulation of the Thalamus for Tremor (1)**

- Tremor suppression was total or clinically significant in 82–91% of operated sides in 179 patients who underwent implantation of thalamic stimulation devices. Results were durable for up to 8 years, and side effects of stimulation were reported as mild and largely reversible.
- These results are at least as good as those associated with thalamotomy. An additional benefit of deep brain stimulation is that recurrence of tremor may be managed by changes in stimulation parameters.

#### **Unilateral or Bilateral Stimulation of the Globus Pallidus or Subthalamic Nucleus for Parkinson's disease Symptoms (2)**

- A wide variety of studies consistently demonstrate that DBS of the globus pallidus or subthalamic nucleus results in significant improvements, as measured by standardized rating scales of neurologic function. The most frequently observed improvements consist of increased waking hours spent in a state of mobility without dyskinesia, improved motor function during “off” periods when levodopa is not effective, reduction in frequency and severity of levodopa-induced dyskinesia during periods when levodopa is working (“on” periods), improvement in cardinal symptoms of PD during periods when medication is not working, and in the case of bilateral DBS of the subthalamic nucleus, reduction in the required daily dosage of levodopa and/or its equivalents. The magnitude of these changes is both statistically significant and clinically meaningful.
- The beneficial treatment effect lasts at least for the 6–12 months observed in most trials. While there is not a great deal of long-term follow-up, the available data are generally positive.
- Adverse effects and morbidity are similar to those known to occur with thalamic stimulation.
- DBS possesses advantages to other treatment options. In comparison to pallidotomy, DBS can be performed bilaterally. The procedure is non-ablative and reversible.

Articles published since these two assessments continue to report positive outcomes for DBS for tremor and PD. In addition, periodic updates of the literature have identified reports on the use of DBS for a variety of neurologic and psychiatric conditions. Following is a summary of key studies.

### **Stimulation of the Thalamus for Essential Tremor and Tremor in Parkinson’s Disease**

In 2008, Schuurman and colleagues reported 5-year follow-up of 65 patients comparing thalamic stimulation and thalamotomy for treatment of tremor due to PD (45 patients), essential tremor (ET) (13 patients), and multiple sclerosis (MS) (10 patients). (3) After 5 years, 48 patients were available for follow-up: 32 with PD, 10 with ET, and 6 with MS. The primary outcome measure was functional status on the Frenchay Activities Index (FAI); secondary measures were tremor severity, frequency of complications, and patients’ assessment of outcome. The mean difference in FAI scores was 4.4 (95% confidence interval [CI]: 1.1–7.7) after 6 months, 3.3 (95% CI: -0.03–6.6) after 2 years, and 4.0 (95% CI: 0.3–7.7) after 5 years in favor of stimulation. Tremor suppression was equally effective after both procedures, and stable in PD patients. A diminished effect was observed in half of the patients with ET and MS. Neurologic adverse effects were higher after thalamotomy. Subjective assessments favored stimulation. Hariz et al. evaluated outcomes of thalamic DBS in patients with tremor-predominant PD who participated in a multicenter European study and reported that, at 6 years post-surgery, tremor was still effectively controlled and appendicular rigidity and akinesia remained stable when compared with baseline. (4)

**Bilateral Stimulation of the Thalamus:** In 2005, Putzke and colleagues reported on a series of 25 patients with ET treated with bilateral DBS for management of midline tremor (head, voice, tongue, trunk). (5) Three patients died of unrelated causes, 1 patient was lost to follow-up due to transfer of care, and 1 patient did not have baseline evaluation; these patients were not included in the analysis. Patients were evaluated at baseline (before implantation of second stimulator), and at 1, 3, 6, 12, 24, and 36 months. At 12 months, evaluations were obtained from 76% of patients; at 36 months, 50% of patients were evaluated. The most consistent improvement on the tremor rating scale during both unilateral and bilateral stimulation was found for head and voice tremor. The incremental improvement over unilateral stimulation through the first 12 months of bilateral stimulation was significant ( $p<0.01$ ). Bilateral stimulation at months 3 and 12 was significantly better than unilateral stimulation at month 3 ( $p<0.05$ ). Small sample size limited analysis at months 24 and 36. Dysarthria was reported in 6 (27%) patients and disequilibrium in 5 patients after bilateral stimulation in staged implantations. No patient reported dysarthria and 2 reported disequilibrium before bilateral stimulation. In 2006, Pahwa et al. reported on long-term follow-up of 45 patients who underwent thalamic DBS, 26 of whom had ET; 18 patients with ET had unilateral and 8 had bilateral implantation. (6) Sixteen patients with unilateral and 7 with bilateral stimulators completed at least part of the 5-year follow-up evaluations. Patients with bilateral stimulation had a 78% improvement in mean motor tremor scores in the stimulation on state compared with baseline at 5-year follow-up ( $p=0.02$ ) and 36% improvement in activity of daily living (ADL) scores. Unilateral stimulation patients improved 46% on motor tremor scores and 51% on ADLs ( $p<0.01$ ). Stimulation-related adverse events were reported in more than 10% of patients with unilateral

and bilateral thalamic stimulators. Most were mild and were reduced with changes in stimulation parameters. Adverse events in patients with bilateral stimulation, such as dysarthria and other speech difficulties, disequilibrium or balance difficulties, and abnormal gait, persisted despite optimization of the stimulation parameters.

### **Stimulation of the Globus Pallidus and Subthalamic Nucleus for Advanced Parkinson's Disease**

A 2006 systematic review of 34 studies (921 patients) examined outcomes following subthalamic stimulation for patients with PD who had failed medical management (e.g., motor fluctuations, dyskinesia, and other medication side effects). (7) Twenty studies, primarily class IV (uncontrolled cohorts or case series), were included in the meta-analysis. Subthalamic stimulation was found to improve ADL by 50% over baseline, as measured by the Unified Parkinson's Disease Rating Scale (UPDRS) part II (decrease of 13.35 points out of 52). There was a 28-point decrease in the UPDRS III score (out of 108), indicating a 52% improvement in the severity of motor symptoms while the patient was not taking medication. A strong relationship was found between the pre-operative dose response to L-dopa and improvements in both the UPDRS II and III. The analysis found a 56% reduction in medication use, a 69% reduction in dyskinesia, and a 35% improvement in quality of life with subthalamic stimulation.

In 2006, the German Parkinson Study Group reported a trial of 78 patient pairs with advanced PD and severe motor symptoms randomized to either subthalamic stimulation or medical management. (8) Subthalamic stimulation improved severity of symptoms without medication in 55 of 78 pairs (from 48 to 28 on the UPDRS III). Improvements in quality of life were greater than medical management in 50 of 78 pairs (average change from 42 to 32 on the 100-point Parkinson's Disease Questionnaire), with 24% to 38% improvements in subscales for mobility, activities of daily living, emotional well-being, stigma, and bodily discomfort. Serious adverse events were more common with neurostimulation (13% vs. 4%) and included a fatal intracerebral hemorrhage. Witt et al. performed an ancillary protocol as part of this multicenter randomized, controlled trial (RCT) to assess neuropsychiatric consequences of DBS in patients with PD. (9) One hundred-twenty-three patients with PD and motor fluctuations who were randomized to DBS or best medical treatment were included in the study. Neuropsychological and psychiatric examinations at baseline and 6 months post-implantation were compared. DBS of the subthalamic nucleus did not reduce overall cognition or affectivity. There was a selective decrease in frontal cognitive functions and an improvement in anxiety in patients after treatment that did not affect improvements in quality of life.

Weaver and colleagues reported 6-month outcomes of a multicenter RCT comparing DBS with best medical therapy for patients with advanced PD in 2009. (10) Of 278 patients who were screened, 255 were randomized; 134 to best medical therapy and 121 to DBS (61 to stimulation of the globus pallidus and 60 to stimulation of the subthalamic nucleus). By intention-to-treat analysis, patients who received DBS gained a mean of 4.6 hours/day of "on" time without troubling dyskinesia compared to no hours gained for patients receiving best medical therapy ( $p<0.001$ ). Seventy-one percent of DBS patients experienced clinically meaningful motor function improvements (i.e.,  $>5$  point change in UPDRS of motor function) versus 32% of best medical therapy group. Significantly greater improvements in quality-of-life measures were achieved by DBS patients. At least one serious adverse event occurred in 49 DBS patients versus 15 in the best medical therapy patients, including 39 related to the surgical procedure and one death secondary to cerebral hemorrhage.

In 2010, Williams et al. reported results from an ongoing randomized, multicenter open-label trial (PD SURG) from 13 neurosurgical centers in the United Kingdom. (11) Included in the study were 366 patients with PD that was not adequately controlled by medical therapy. Patients were randomized to surgery (all had DBS) and best medical therapy, or to best medical therapy alone. The study was designed to detect a 10-point difference (regarded as clinically important) in the Parkinson's disease questionnaire (PDQ) summary index. Five of 183 patients randomized to surgery did not have surgery, and 12 of 183 patients randomized to medical therapy had surgery within the first year of the study (patients were analyzed in the treatment group to which they were randomized). In 174 patients, the subthalamic nucleus was the surgical target, and 176 of 178 procedures were bilateral. At 1 year, the

mean improvement in the primary outcome measure, the PDQ summary index, was 5.0 points in the DBS group and 0.3 points in the control group. The difference in mean change in PDQ between the 2 groups was -8.9 for the mobility domain, -12.4 for the daily living domain, and -7.5 for the bodily discomfort domain. Differences between groups in the other domains were not significant. Thirty-six (19%) patients had serious surgery-related adverse events; there was one procedure-related death. The most common surgery-related serious adverse events were infections (n=16).

Another European multicenter study assessed whether subthalamic stimulation might maintain quality of life and motor function if performed earlier in the course of the disease. (12) Ten matched patient pairs younger than 55 years of age with mild to moderate motor signs were randomly assigned to DBS or medical management. There was no difference in the severity of parkinsonian motor disability while receiving medication. However, in the medically treated patients, both the daily dose of levodopa and the severity of levodopa-induced motor complications increased over the 18 months of the study (12% and 15%, respectively), while in the surgical patients, the daily dose of levodopa was reduced by 57%, and the severity of levodopa-induced motor complications improved by 83%. Additional studies are needed to determine the long-term effect of subthalamic stimulation in this younger patient population.

Appleby et al. reported a meta-analysis of adverse events associated with DBS in order to assess the risks and benefits of the treatment as they relate to its potential use in the psychiatric setting in 2007. (13) They concluded that DBS is an effective treatment for PD, dystonia, and ET, and rates of depression, cognitive impairment, mania, and behavior change are low. Prevalence of depression was 2–4%, mania 0.9–1.7%, emotional changes 0.1–0.2%, and suicidal ideation/suicide attempt was 0.3–0.7%. The completed suicide rate was 0.16–0.32%. In light of the rate of suicide in patients treated with DBS, particularly with thalamic and globus pallidus stimulation, the authors argue for prescreening patients for suicide risk.

### **Deep Brain Stimulation for the Treatment of Dystonia**

DBS for the treatment of primary dystonia received FDA approval through the Humanitarian Device Exemption (HDE) process in 2003. The HDE approval process is available for conditions that affect fewer than 4,000 Americans per year. According to this approval process, the manufacturer is not required to provide definitive evidence of efficacy, but only probable benefit. The approval was based on the results of DBS in 201 patients represented in 34 manuscripts. (14) There were 3 studies that reported at least 10 cases of primary dystonia. In these studies, clinical improvement ranged from 50% to 88%. A total of 21 pediatric patients were studied; 81% were older than 7 years. Among these patients, there was an approximate 60% improvement in clinical scores. As noted in the analysis of risk and probable benefit, the only other treatment options for chronic refractory primary dystonia are neurodestructive procedures. DBS provides a reversible alternative. The FDA Summary of Safety and Probable Benefit states, “Although there are a number of serious adverse events experienced by patients treated with deep brain stimulation, in the absence of therapy, chronic intractable dystonia can be very disabling and, in some cases, progress to a life-threatening stage or constitute a major fixed handicap. When the age of dystonia occurs prior to the individual reaching their full adult size, the disease not only can affect normal psychosocial development but also cause irreparable damage to the skeletal system. As the body of the individual is contorted by the disease, the skeleton may be placed under constant severe stresses that may cause permanent disfigurement. Risks associated with deep brain stimulation for dystonia appear to be similar to the risk associated with the performance of stereotactic surgery and the implantation of deep brain stimulation systems for currently approved indications, except when used in either child or adolescent patient groups.”

Since the FDA approval, there have been additional published trials of deep brain stimulation for dystonia, which continue to report positive results. (15) Vidailhet and colleagues reported the results of a prospective multi-institutional case series of 22 patients with primary generalized dystonia. (16) Symptoms were evaluated prior to surgery and at several points up to 1 year of follow-up, in a double-blind fashion with the stimulator turned on and off. Dystonia scores were significantly better with the neurostimulator turned on. Vidailhet et al. compared outcomes at 3 years with those reported at 1 year for the 22 patients in their study of bilateral, pallidal DBS for generalized dystonia and found that the

motor improvement observed at 1 year was maintained. (16, 17) At 3 years, measures of cognition and mood were unchanged from baseline and 1-year evaluations. Egidi et al. retrospectively reviewed records of 69 patients treated in multiple Italian centers with DBS implanted in the globus pallidus; 37 patients had primary and 32 had secondary dystonia. Improvement of at least 50% in Burke-Fahn-Marsden severity scale was reached by 45% of primary and 37% of secondary dystonia patients at 3–84 months' follow-up (longer than 24 months in half of the patients). (18)

In 2006, the Deep-Brain Stimulation for Dystonia Study Group compared bilateral pallidal neurostimulation with sham stimulation in 40 patients with dystonia who had failed medical management (3-month randomized trial with a 6-month open-label extension). (19) Blinded assessment with the Burke-Fahn-Marsden Dystonia Rating Scale (BFMDRS) found improvements in the movement score (16 points vs. 1.6 points in sham controls), which corresponded to a 39% reduction in symptoms. Disability scores improved by 4 points in the neurostimulation group compared with a 0.8-point improvement in the control subjects (38% improvement). The study found a 30% improvement in quality of life (change of 10 vs. 4 points in controls) following stimulation of the globus pallidus. There was high variability in baseline scores and in the magnitude of improvement; 6 patients (17%) were considered to have failed treatment (<25% improvement), 5 patients (25%) improved by more than 75%. No single factor was found to predict the response to treatment. Independent assessors found similar improvements in the control group after the 6-month open-label extension. Thirty-eight patients (95%) agreed to be followed up annually, and 80% of patients completed the 5-year follow-up. (20) Intention-to-treat analysis showed significant improvements in dystonia severity at 6 months (-47.9%), 3 years (-61.1%), and 5 years (-57.8% compared with baseline). The unmasked raters tended to score dystonia severity as higher than the 2 masked raters in the original study. Responder analysis (>25% on the BFMDRS) indicated a positive response in 83% of 36 patients at 6 months, 94% of 31 patients at 3 years, and 81% of 32 patients at 5 years. There were 21 serious adverse events that required hospitalization. Almost all serious adverse events were device-related including subcutaneous infection, lead dislodgement/lead breakage, and stimulator malfunction. The most common non-serious adverse event was dysarthria.

**Stimulation of the Globus Pallidus for the Treatment of Tardive Dyskinesia and Tardive Dystonia**  
Stimulation of the globus pallidus has been examined as a treatment of tardive dyskinesia in a Phase II double-blinded (presence and absence of stimulation) multicenter study. (21) The trial was stopped early due to successful treatment (greater than 40% improvement) in the first 10 patients.

Outcomes on motor function, quality of life, and mood in a series 9 patients treated with DBS of the globus pallidus internus for tardive dystonia were reported by Gruber et al. in 2009. (22) One week and 3 to 6 months after surgery, BFMDRS motor scores were improved by 56.4 +/- 26.7% and 74.1 +/- 15.8%, BFMDRS disability scores by 62.5 +/- 21% and 88.9 +/- 10.3%, and Abnormal Involuntary Movement Scale (AIMS) scores by 52.3 +/- 24.1% and 69.5 +/- 27.6%, respectively. At last follow-up (mean 41 months, range 18- 90 months), BFMDRS motor scores were reduced compared to presurgical assessment by 83 +/- 12.2%, BFMDRS disability score by 67.7 +/- 28%, and AIMS scores by 78.7 +/- 19.9%.

## **Epilepsy**

In 2010, Fisher et al. reported a U.S. multicenter, double-blind, randomized trial of bilateral stimulation of the anterior nuclei of the thalamus for epilepsy (SANTE). (23) Included were 110 patients 18- to 65-years-old, with partial seizures including secondarily generalized seizures, at least 6 per month, but no more than 10 per day. An additional 47 patients were enrolled in the study but did not undergo implantation. At least 3 antiepileptic drugs must have failed to produce adequate seizure control prior to baseline, with 1 to 4 antiepileptic drugs used at the time of study entry. Half of the patients were randomized to stimulation during a 3-month blinded phase; then all patients received unblinded stimulation. The baseline monthly median seizure frequency was 19.5. During the first and second months of the blinded phase, the difference in seizure reduction between stimulation on and stimulation off (-42.1% vs. -28.7%, respectively) was not significantly different. In the last month of the blinded phase, the stimulated group had a greater reduction in seizures compared with the control group (-

40.4% vs. -14.5% in controls). The median change in seizure frequency was -41% at 13 months and -56% at 25 months. The stimulation group experienced fewer seizure-related injuries than patients in the control group (7.4% vs. 25.5%, respectively). Cognition and mood showed no group differences, but participants in the stimulated group were more likely to report depression (8 vs. 1) or memory problems (7 vs. 1 – both respectively) as adverse events. There was a progressive reduction in seizure frequency over long-term follow-up. By 2 years, 54% of patients had a seizure reduction of equal to or greater than 50%, and 14 patients (13%) were seizure-free for at least 6 months. The most common device-related adverse events were paresthesias in 18.2% of participants, implant site pain in 10.9%, and implant site infection in 9.1%. Eighteen participants (16.4%) withdrew from the study after the implantation because of adverse events. There were 5 deaths, none of which were considered to be device-related. Although some patients appear to have benefited from treatment during the extended follow-up phase, the difference between groups in the blinded portion of the study was modest. Additional study is needed to establish the safety and efficacy of this treatment.

### **Huntington's Disease**

No controlled trials of DBS for Huntington's disease were identified.

### **Multiple Sclerosis**

In 2008, Schuurman and colleagues reported 5-year follow-up of 65 patients comparing thalamic stimulation and thalamotomy for treatment of tremor due to PD (45 patients), ET (13 patients), and MS (10 patients). (3) After 5 years, 48 patients were available for follow-up: 32 with PD, 10 with ET, and 6 with MS. The primary outcome measure was functional status on the FAI; secondary measures were tremor severity, frequency of complications, and patients' assessment of outcome. The mean difference in FAI scores was 4.4 (95% CI: 1.1–7.7) after 6 months, 3.3 (95% CI: -0.03–6.6) after 2 years, and 4.0 (95% CI: 0.3–7.7) after 5 years in favor of stimulation. Tremor suppression was equally effective after both procedures, and stable in PD patients. A diminished effect was observed in half of the patients with ET and MS. Small numbers of patients with MS limit conclusions with respect to this condition.

### **Tourette Syndrome**

A 2012 systematic review identified 25 published studies, representing data from 69 patients that reported on the efficacy of DBS in the treatment of Tourette syndrome. (24) However, only 3 studies with methodologic quality ratings of fair to poor met the inclusion criteria for evidence-based analysis. These 3 studies are described below. The authors recommend that DBS continues to be considered an experimental treatment for severe, medically refractory tics.

Another systematic review from 2012 examined patient and target selection for DBS of Tourette syndrome. (25) The majority of clinical trials for DBS in Tourette syndrome have targeted the medial thalamus at the crosspoint of the centromedian nucleus, substantia periventricularis, and nucleus ventro-oralis internus. Other targets that have been investigated include the subthalamic nucleus, caudate nucleus, globus pallidus internus, and the anterior limb of the internal capsule and nucleus accumbens. The review found no clear consensus in the literature for which patients should be treated and what the best target is. Additional study is needed to clarify these issues.

Three small cross-over studies of DBS for Tourette syndrome have been identified. One compared unilateral and bilateral thalamic stimulation (5 patients) and the other (3 patients) compared thalamic, pallidal, simultaneous thalamic and pallidal, and sham stimulation. (26, 27) The best improvements were found with the ventromedial pallidal stimulation.

In 2011, Ackermans et al. reported preliminary results of a double-blind crossover trial of thalamic stimulation in 6 patients with refractory Tourette syndrome. (28) Tic severity during 3 months of stimulation was significantly lower than during the 3 months with the stimulator turned off, with a 37% improvement on the Yale Global Tic Severity Scale (mean 25.6 vs. 41.1) and a decrease in tic severity of 49% at 1 year after surgery compared to preoperative assessments (mean 21.5 vs. 42.2 – both respectively). Secondary outcomes (change in associated behavioral disorder and mood) were not altered by the stimulation. Serious adverse events included one small hemorrhage ventral to the tip of

the electrode, one infection of the pulse generator, subjective gaze disturbances, and reduction of energy levels in all patients. The interim analysis led to the termination of the trial. The authors commented that further RCTs on other targets are urgently needed, since the search for the optimal one is still ongoing.

### **Hypothalamic Stimulation for the Treatment of Cluster Headaches and Facial Pain**

Deep brain stimulation of the posterior hypothalamus for the treatment of chronic cluster headaches has been investigated, since functional studies have suggested cluster headaches have a central hypothalamic pathogenesis.

In 2010, Fontaine et al. published results from a prospective crossover, double-blind, multicenter study in 11 patients with DBS of the posterior hypothalamus for severe refractory chronic cluster headache. (29) The randomized phase compared active and sham stimulation during 1-month periods, and was followed by a 1-year open phase. Severity of cluster headache was assessed by the weekly attacks frequency (primary outcome), pain intensity, sumatriptan injections, emotional impact, and quality of life (short-form 12 [SF-12]). During the randomized phase, no significant change in primary and secondary outcome measures was observed between active and sham stimulation. At the end of the open phase, 6 of 11 patients reported a greater than 50% reduction in the weekly frequency of attacks.

Another research group from Europe has published several case series (potentially overlapping) on DBS of the ipsilateral posterior hypothalamus in patients with chronic cluster headache. (30, 31) Stimulation was reported to result in long-term pain relief (1–26 months of follow-up) without significant adverse effects in 16 patients with chronic cluster headaches and in 1 patient with neuralgiform headache; treatment failed in 3 of 3 patients who had atypical facial pain. Controlled studies are needed to evaluate the long-term safety and effectiveness of DBS for chronic cluster headaches.

### **Treatment-Resistant Depression**

A variety of target areas are being investigated for DBS of treatment-resistant depression. A systematic review from 2014 identified 22 published reports with 6 different approaches/targets including the nucleus accumbens, ventral striatum/ventral capsule, subgenual cingulate cortex, lateral habenula, inferior thalamic nucleus, and medial forebrain bundle. (32) Only 3 of the studies identified were controlled with sham stimulation periods, and as of December 2013, there were 2 unpublished multicenter, randomized, controlled trials evaluating subgenual cingulate cortex and ventral striatum/ventral capsule DBS that had been terminated due to futility (interim analysis demonstrating very low probability of success if trial was completed as planned).

A representative within-subject study is described in a 2012 report by Holtzheimer et al, who reported a Phase I/II open-label trial of DBS of the subcallosal cingulate white matter with a single-blind sham lead-in phase for treatment-resistant unipolar and bipolar depression. (33) Ten patients with treatment-resistant major depressive disorder and 7 patients with treatment-resistant bipolar II disorder were included in the study. Inclusion criteria included a current major depressive episode of at least 12 months' duration, a score of 20 or higher on the Hamilton Depression Rating Scale (HAM-D), not responding to at least 4 adequate antidepressant treatments, and a lifetime failure or inability to receive electroconvulsive therapy. The mean HAM-D score was 20.5 at the end of the sham lead-in phase, decreasing to 13.1 at 24 weeks (n=16), 13.6 at 1 year (n=14), and 7.3 at 2 years (n=11). Remission rates, defined as a HAM-D score of less than 8, were 18% at 24 weeks, 36% at 1 year, and 58% after 2 years. Response rates, defined as 50% or greater change in the HAM-D, were 41% after 24 weeks, 36% after 1 year and 92% after 2 years. The first 3 patients underwent a single-blind discontinuation phase after 24 weeks, and all 3 had full relapse with increased suicidal ideation. Because of patient safety concerns, this phase was eliminated for subsequent patients. No patient achieving remission experienced a relapse during stimulation. Efficacy was similar for patients with major depression and those with bipolar depression.

### **Obsessive-Compulsive Disorder**

A systematic review of DBS for obsessive-compulsive disorder (OCD) was published in 2011 by de Koning et al. (34) The review included 9 case studies and 7 controlled studies with a blinded on-off phase. It was estimated from the published trials and case studies that more than 100 individuals have received experimental DBS for OCD in 5 different targets. These targets are the anterior limb of the internal capsule (ALIC), subthalamic nucleus, ventral capsule/ventral striatum, nucleus accumbens, and inferior thalamic peduncle. The most common measure of efficacy is a reduction in the Yale-Brown Obsessive Compulsive Scale (Y-BOCS). The Y-BOCS is a 10-item scale in which higher scores reflect more intense symptoms, and a score of 24 or more (of a possible 40) is considered severe illness. Most studies designate a therapeutic response as a Y-BOCS reduction of 35% or more from the pretreatment baseline, with a reduction of 25% or more considered a partial response.

Subthalamic Nucleus: A crossover, double-blind, multicenter study of DBS of the subthalamic nucleus for treatment of refractory OCD was reported by Mallet et al. in 2008. (35) Eighteen patients were enrolled, one withdrew and one required removal of the stimulator before randomization because of infection. Three months after surgery, 8 patients were randomly assigned to receive active stimulation for 3 months, followed by 1 month of washout, then 3 months of sham stimulation (on-off group). The other group followed the same treatment schedule in reverse (off-on group). New or worsening symptoms were classified as adverse events. Medication was held constant during the 10-month protocol, except for transient increase in benzodiazepine therapy in 3 patients and augmentation of neuroleptic treatment in one patient for exacerbated anxiety. The Y-BOCS score was significantly lower at the end of active stimulation than at the end of the sham stimulation (mean score, 19 +/- 8 vs. 28 +/- 7; p=0.01) independent of the group and the period. No significant carryover effect between treatment phases was detected. Patients who had active stimulation first (on-off group) tended to have a larger treatment effect than the off-on group (p=0.06).

Outcomes on secondary measures of global health and functioning were significantly better at the end of the stimulation period. Scores on Montgomery and Asberg Depression Scale (MADRS), Brief Scale for Anxiety, neuropsychological ratings, and self-reported disability (Sheehan Disability Scale) did not differ significantly at the end of treatment and sham sessions. Fifteen serious adverse events were reported in 11 patients, the most serious a parenchymal brain hemorrhage. Transient motor and psychiatric symptoms induced by active stimulation resolved spontaneously or with adjustment of stimulation settings. Seven behavioral adverse events were reported in 5 patients during stimulation. Hypomania was the main psychiatric serious adverse event; symptoms resolved with adjustment of stimulation settings. The authors note that the multicenter design might be a limitation of the study because of variation in targeting of stimulation. In addition, in order to preserve blinding, stimulation settings were kept below the threshold known to induce adverse effects and may have been too low to reduce symptoms. They conclude that their finding suggests that DBS may lessen severity of symptoms; however, serious adverse events did occur. Larger studies with longer follow-up are needed including evaluation of quality of life and ability to function in social and work situations.

Nucleus Accumbens: Denys et al. reported a double-blind crossover study of bilateral DBS of the nucleus accumbens in 16 patients with refractory OCD in 2010. (36) Patients with a score of equal to or greater than 28 on the Y-BOCS, and an equal to or greater than 5-year history of OCD that was refractory to medical treatment were included. Out of 101 patients screened for the study, 16 underwent implantation. The study consisted of an open 8-month treatment phase, followed by a double-blind crossover phase with randomly assigned 2-week periods of active or sham stimulation, ending with an open 12-month maintenance phase. Once a decrease in Y-BOCS was obtained, a standardized cognitive behavioral therapy (CBT) program was added. In the open phase, the mean Y-BOCS score decreased by 46% from 33.7 at baseline to 18.0 after 8 months. Nine of 16 patients were responders (Y-BOCS decrease >35%), with a mean decrease of 23.7 points. In the double-blind, sham-controlled phase (n=14), there was a mean 8.3 point difference in the Y-BOCS score between active and sham stimulation. Depression and anxiety decreased significantly, with a mean difference in Hamilton anxiety (HAM-A) scores of 12.1 and in Hamilton depression (HAM-D) scores of 11.3. Except for mild forgetfulness and word-finding problems, no permanent adverse events were reported. The most prominent adverse event related to stimulation was elevated mood or hypomania.

A double-blind crossover study of unilateral DBS of the nucleus accumbens was reported by Huff et al. in 2010. (37) Patients with a score of equal to or greater than 25 on the Y-BOCS, and an equal to or greater than 5-year history of OCD that was refractory to medical treatment were included. Ten patients received 3 months of DBS followed by 3 months of sham stimulation, or vice versa. After 6 months, stimulation was continued unblinded with the option to change stimulation parameters every 3 months (including activation of electrodes in the ALIC). The patients had an examination at baseline, within the first week, and at 3, 6, 9, and 12 months by a psychiatrist who was blinded to the treatment condition. The mean Y-BOCS at baseline was 32.2. There was no difference in Y-BOCS during the crossover period with a score of 27.9 during the on period and 31.1 during the off period. After 12 months the Y-BOCS had significantly decreased to 25.4. Logistic regression revealed no independent effect for changes in stimulation amplitude, changes in active contacts, or changes in medication. Five of 10 patients showed a decrease of equal to or greater than 25%, indicating a partial response. Only 1 patient showed a decrease in Y-BOCS of greater than 35%. Depression, global functioning, and quality of life improved within 1 year, while anxiety, global symptom severity, and cognitive function showed no significant changes.

**Ventral Capsule/Ventral Striatum:** Goodman et al. reported a double-blinded pilot crossover study of DBS of the ventral capsule/ventral striatum in 6 patients. (38) Patients with a score of equal to or greater than 28 on the Y-BOCS and a 5-year or longer history of OCD that was refractory to medical treatment were included. All 6 patients had a lifetime diagnoses of major depression that was deemed secondary to OCD, one met criteria for a current diagnosis of major depression. The mean duration of illness was 24 years (range, 11-35 years). The first patient was implanted in 2003; the sixth patient completed 12 months of DBS in 2008. The baseline Y-BOCS was 33.7. For the crossover phase, there was a reduction of 5.33 points with the stimulator turned on (n=3) and -0.67 with the stimulator off (n=3, not significantly different). After 12 months of stimulation, 4 (66.7%) of patients were responders (>35% improvement and a score <16 on the Y-BOCS). Depressive symptoms improved significantly in the group as a whole; global functioning improved in the 4 responders. The authors concluded that future research should attend to subject selection, lead location, DBS programming, and mechanisms underlying the therapeutic benefits.

**Section Summary:** The literature on DBS for OCD consists of a small number of randomized, double-blind crossover trials. These trials report improvements in OCD measures associated with DBS, but the differences were not statistically significant in all trials. This literature also indicates that the optimal stimulation site has not been determined. One small study of stimulation in the subthalamic nucleus showed short-term improvement in the Y-BOCS score, while stimulation of the nucleus accumbens led to conflicting results in 2 small cross-over studies. A pilot study of stimulation in the ventral striatum in 6 patients suggests that this might be an effective area of stimulation for some patients, although the mean improvement in Y-BOCS was modest. Additional research with a larger number of patients is needed to evaluate the effect of DBS for treatment-resistant OCD, including determination of the most effective site of stimulation.

## **Other**

The evidence on deep brain stimulation for anorexia nervosa, alcohol addiction, Alzheimer Disease and chronic pain consists of review articles or small case series. These are not adequate to make a determination of efficacy.

## **Ongoing and Unpublished Clinical Trials**

A search of the online site [www.clinicaltrials.gov](http://www.clinicaltrials.gov) in June 2014 identified over 100 active and unpublished trials on DBS for a variety of neurological conditions. Of particular note is INTREPID (NCT01839396), a manufacturer-sponsored multi-center, double-blinded, randomized controlled trial to assess the safety and effectiveness of the Vercise™ DBS System in patients with Parkinson disease. All patients will be implanted with the DBS stimulator; the control group will receive low dose stimulation for the first 12 weeks of the study and the active group will receive moderate dose stimulation. The manufacturer expects to enroll 310 patients with study completion anticipated July 2021.

## Summary

Deep brain stimulation (DBS) has been shown to be effective for the treatment of tremor, advanced Parkinson's disease, and dystonia. Evidence for efficacy of DBS for Tourette syndrome, treatment-resistant depression, obsessive-compulsive disorder, Alzheimer disease, anorexia nervosa, alcohol addiction, chronic pain, epilepsy, tardive dystonia, and cluster headache is based on experience with small numbers of patients. In addition, the appropriate candidates and most effective target areas for DBS are under investigation. Additional controlled studies with a larger number of subjects are required to evaluate the role of DBS for these conditions.

## Practice Guidelines and Position Statements

The American Academy of Neurology (AAN) published an updated guideline on the treatment of essential tremor in 2011. (39) There were no changes from the conclusions and recommendations of the 2005 practice parameters regarding DBS for essential tremor. (40) The guidelines stated that bilateral DBS of the thalamic nucleus may be used to treat medically refractory limb tremor in both upper limbs (level C, possibly effective) but that there were insufficient data regarding the risk:benefit ratio of bilateral versus unilateral DBS in the treatment of limb tremor. There was insufficient evidence to make recommendations regarding the use of thalamic DBS for head or voice tremor (level U, treatment is unproven).

2006 Guidelines from AAN on the treatment of PD with motor fluctuations and dyskinesia found that although the criteria are evolving, patients with PD who are considered candidates for DBS include levodopa-responsive, non-demented, and neuropsychiatrically intact patients who have intractable motor fluctuations, dyskinesia, or tremor. (41) AAN concluded that DBS of the subthalamic nucleus may be considered as a treatment option in PD patients to improve motor function and to reduce motor fluctuations, dyskinesia, and medication usage (Level C – possibly effective), but found insufficient evidence to make any recommendations about the effectiveness of DBS of the globus pallidus or the ventral intermediate nucleus of the thalamus in reducing motor complications or medication usage, or in improving motor function in PD patients.

2010 Guidelines from AAN on the treatment of non-motor symptoms of Parkinson disease found insufficient evidence for the treatment of urinary incontinence with DBS of the subthalamic nucleus. (42) AAN found that DBS of the subthalamic nucleus possibly improves sleep quality in patients with advanced Parkinson disease. However, none of the studies performed DBS to treat insomnia as a primary symptom, and DBS of the subthalamic nucleus is not currently used to treat sleep disorders.

2013 Guidelines from AAN on the treatment of tardive syndromes states that the available evidence, which consists of Class IV studies comprising case reports or small case series, is insufficient to support or refute pallidal DBS for tardive syndromes. (43)

The American Psychiatric Association states in a 2007 guideline on treatment of patients with obsessive-compulsive disorder that DBS may be recommended on the basis of individual circumstances. (44) DBS is not mentioned in the 2010 guideline on treatment of major depressive disorder or in the 2013 Guideline Watch.

The Canadian Network for Mood and Anxiety Treatments (CANMAT) 2009 clinical guideline for management of major depressive disorder in adults states that there is emerging evidence to support DBS as an experimental intervention for patients with treatment-refractory depression. (45) There is no consensus on the most effective target brain region for implantation, although 3 regions have been explored (subcallosal cingulated gyrus, the nucleus accumbens, and the ventral caudate/ventral striatum region).

The European Society for the Study of Tourette Syndrome (ESSST) published guidelines on DBS in 2011. (46) The guidelines state that DBS for Tourette syndrome is still in its infancy and that there are

no randomized controlled studies available that include a sufficiently large number of patients. There was general agreement among the workgroup members that DBS should only be used in adult, treatment-resistant, and severely affected patients, and it was highly recommended that DBS be performed in the context of controlled and double-blind trials including larger and carefully characterized groups of patients.

The U.K.'s National Institute for Health and Care Excellence (NICE, previously the National Institute for Clinical Excellence) has published Interventional Procedure Guidance documents on deep brain stimulation.

- Tremor and Dystonia: In 2006, NICE made the same statement for use of DBS for treatment of tremor and dystonia. (47) Unilateral and bilateral stimulation of structures responsible for modifying movements, such as the thalamus, the globus pallidus, and the subthalamic nucleus, which interact functionally with the substantia nigra, are included in both guidance statements.
- Refractory Chronic Pain Syndromes (excluding headache): 2011 guidance states that there is evidence that DBS is efficacious in some patients who are refractory to other forms of pain control and that this procedure may be used provided that normal arrangements are in place for clinical governance, consent, and audit. (48) Patients should be informed that DBS may not control their chronic pain symptoms and that possible risks associated with this procedure include the small risk of death.
- Intractable Trigeminal Autonomic Cephalgias: 2011 guidance states that current evidence on the efficacy of DBS for intractable trigeminal autonomic cephalgias (e.g., cluster headaches) is limited and inconsistent, and the evidence on safety shows that there are serious but well-known side effects. (49) Therefore this procedure should only be used with special arrangements for clinical governance, consent, and audit or research.
- Refractory Epilepsy: 2012 guidance states that the evidence on the efficacy of DBS for refractory epilepsy is limited in both quantity and quality. (50) The evidence on safety shows that there are serious but well-known side effects. Therefore, this procedure should only be used with special arrangements for clinical governance, consent, and audit or research.
- Parkinson's Disease: In 2003, NICE stated that current evidence on the safety and efficacy of DBS for treatment of Parkinson's disease appears adequate to support the use of the procedure, provided that normal arrangements are in place for consent, audit, and clinical governance. (51)
- Parkinson Disease: In 2006, NICE published a clinical guideline on the diagnosis and management of Parkinson disease in primary and secondary care. (52) With the evidence at that time it was not possible to decide if the subthalamic nucleus or globus pallidus interna is the preferred target for deep brain stimulation for people with PD, or whether one form of surgery is more effective or safer than the other. Based on level 3 or 4 evidence, NICE concluded that thalamic deep brain stimulation may be considered as an option in people with PD who predominantly have severe disabling tremor and where subthalamic nucleus stimulation cannot be performed

## **U.S. Preventative Services Task Force (USPSTF) Recommendations**

Deep brain stimulation is not a preventative service.

## **Medicare National Coverage**

Effective for services furnished on or after April 1, 2003, Medicare will cover unilateral or bilateral thalamic ventralis intermedius nucleus (VIM) deep brain stimulation (DBS) for the treatment of essential tremor (ET) and/or parkinsonian tremor and unilateral or bilateral subthalamic nucleus (STN) or globus pallidus interna (GPI) DBS for the treatment of Parkinson's disease (PD) when the following conditions are met. (53)

1. DBS devices must be U.S. Food and Drug Administration (FDA) approved devices for DBS or devices used in accordance with FDA-approved protocols governing Category B Investigational Device Exemption (IDE) DBS clinical trials.

2. For thalamic VIM DBS, patients must meet all of the following criteria:
  - a) Diagnosis of ET based on postural or kinetic tremors of hand(s) without other neurologic signs, or diagnosis of idiopathic PD (presence of at least 2 cardinal PD features (tremor, rigidity or bradykinesia)) which is of a tremor-dominant form.
  - b) Marked disabling tremor of at least level 3 or 4 on the Fahn-Tolosa-Marin Clinical Tremor Rating Scale (or equivalent scale) in the extremity intended for treatment, causing significant limitation in daily activities despite optimal medical therapy.
  - c) Willingness and ability to cooperate during conscious operative procedure, as well as during post-surgical evaluations, adjustments of medications and stimulator settings.
3. For STN or GPi DBS, patients must meet all of the following criteria:
  - a) Diagnosis of PD based on the presence of at least 2 cardinal PD features (tremor, rigidity or bradykinesia).
  - b) Advanced idiopathic PD as determined by the use of Hoehn and Yahr stage or Unified Parkinson's Disease Rating Scale (UPDRS) part III motor subscale.
  - c) L-dopa responsive with clearly defined "on" periods.
  - d) Persistent disabling Parkinson's symptoms or drug side effects (e.g., dyskinesias, motor fluctuations, or disabling "off" periods) despite optimal medical therapy.
  - e) Willingness and ability to cooperate during conscious operative procedure, as well as during post-surgical evaluations, adjustments of medications and stimulator settings

DBS is not covered for ET or PD patients with any of the following:

1. Non-idiopathic Parkinson's disease or "Parkinson's Plus" syndromes.
2. Cognitive impairment, dementia or depression, which would be worsened by or would interfere with the patient's ability to benefit from DBS.
3. Current psychosis, alcohol abuse or other drug abuse.
4. Structural lesions such as basal ganglionic stroke, tumor or vascular malformation as etiology of the movement disorder.
5. Previous movement disorder surgery within the affected basal ganglion.
6. Significant medical, surgical, neurologic or orthopedic co-morbidities contraindicating DBS surgery or stimulation.

## References

1. Blue Cross and Blue Shield Technology Evaluation Center. Deep brain stimulation of the thalamus for tremor. TEC Assessment 1997; Volume 12, Tab 20.
2. Blue Cross and Blue Shield Technology Evaluation Center. Bilateral deep brain stimulation of the subthalamic nucleus or the globus pallidus interna for treatment of advanced Parkinson's disease. TEC Assessment 2001; Volume 16, Tab 16.
3. Schuurman PR, Bosch DA, Merkus MP et al. Long-term follow-up of thalamic stimulation versus thalamotomy for tremor suppression. *Mov Disord* 2008; 23(8):1146-53.
4. Hariz MI, Krack P, Alesch F et al. Multicentre European study of thalamic stimulation for parkinsonian tremor: a 6 year follow-up. *J Neurol Neurosurg Psychiatry* 2008; 79(6):694-9.
5. Putzke JD, Uitti RJ, Obwegeser AA et al. Bilateral thalamic deep brain stimulation: midline tremor control. *J Neurol Neurosurg Psychiatry* 2005; 76(5):684-90.
6. Pahwa R, Lyons KE, Wilkinson SB et al. Long-term evaluation of deep brain stimulation of the thalamus. *J Neurosurg* 2006; 104(4):506-12.
7. Kleiner-Fisman G, Herzog J, Fisman DN et al. Subthalamic nucleus deep brain stimulation: summary and meta-analysis of outcomes. *Mov Disord* 2006; 21 Suppl 14:S290-304.
8. Deuschl G, Schade-Brittinger C, Krack P et al. A randomized trial of deep-brain stimulation for Parkinson's disease. *N Engl J Med* 2006; 355(9):896-908.
9. Witt K, Daniels C, Reiff J et al. Neuropsychological and psychiatric changes after deep brain stimulation for Parkinson's disease: a randomised, multicentre study. *Lancet Neurol* 2008; 7(7):605-14.
10. Weaver FM, Follett K, Stern M et al. Bilateral deep brain stimulation vs best medical therapy for patients with advanced Parkinson disease: a randomized controlled trial. *JAMA* 2009; 301(1):63-73.

11. Williams A, Gill S, Varma T et al. Deep brain stimulation plus best medical therapy versus best medical therapy alone for advanced Parkinson's disease (PD SURG trial): a randomised, open-label trial. *Lancet Neurol* 2010; 9(6):581-91.
12. Schupbach WM, Maltete D, Houeto JL et al. Neurosurgery at an earlier stage of Parkinson disease: a randomized, controlled trial. *Neurology* 2007; 68(4):267-71.
13. Appleby BS, Duggan PS, Regenbberg A et al. Psychiatric and neuropsychiatric adverse events associated with deep brain stimulation: A meta-analysis of ten years' experience. *Mov Disord* 2007; 22(12):1722-8.
14. U.S. Food and Drug Administration. Summary of Safety and Probable Benefit. Medtronic Activa Dystonia Therapy. 2003. Available online at: [http://www.accessdata.fda.gov/cdrh\\_docs/pdf2/H020007b.pdf](http://www.accessdata.fda.gov/cdrh_docs/pdf2/H020007b.pdf)
15. Halbig TD, Gruber D, Kopp UA et al. Pallidal stimulation in dystonia: effects on cognition, mood, and quality of life. *J Neurol Neurosurg Psychiatry* 2005; 76(12):1713-6.
16. Vidailhet M, Vercueil L, Houeto JL et al. Bilateral deep-brain stimulation of the globus pallidus in primary generalized dystonia. *N Engl J Med* 2005; 352(5):459-67.
17. Vidailhet M, Vercueil L, Houeto JL et al. Bilateral, pallidal, deep-brain stimulation in primary generalised dystonia: a prospective 3 year follow-up study. *Lancet Neurol* 2007; 6(3):223-9.
18. Egidi M, Franzini A, Marras C et al. A survey of Italian cases of dystonia treated by deep brain stimulation. *J Neurosurg Sci* 2007; 51(4):153-8.
19. Kupsch A, Benecke R, Muller J et al. Pallidal deep-brain stimulation in primary generalized or segmental dystonia. *N Engl J Med* 2006; 355(19):1978-90.
20. Volkmann J, Wolters A, Kupsch A et al. Pallidal deep brain stimulation in patients with primary generalised or segmental dystonia: 5-year follow-up of a randomised trial. *Lancet Neurol* 2012; 11(12):1029-38.
21. Damier P, Thobois S, Witjas T et al. Bilateral deep brain stimulation of the globus pallidus to treat tardive dyskinesia. *Arch Gen Psychiatry* 2007; 64(2):170-6.
22. Gruber D, Trittenberg T, Kivi A et al. Long-term effects of pallidal deep brain stimulation in tardive dystonia. *Neurology* 2009; 73(1):53-8.
23. Fisher R, Salanova V, Witt T et al. Electrical stimulation of the anterior nucleus of thalamus for treatment of refractory epilepsy. *Epilepsia* 2010; 51(5):899-908.
24. Steeves T, McKinlay BD, Gorman D et al. Canadian guidelines for the evidence-based treatment of tic disorders: behavioural therapy, deep brain stimulation, and transcranial magnetic stimulation. *Can J Psychiatry* 2012; 57(3):144-51.
25. Pansaon Piedad JC, Rickards HE, Cavanna AE. What Patients with Gilles de la Tourette Syndrome Should be Treated with Deep Brain Stimulation and What is the Best Target? *Neurosurgery* 2012.
26. Maciunas RJ, Maddux BN, Riley DE et al. Prospective randomized double-blind trial of bilateral thalamic deep brain stimulation in adults with Tourette syndrome. *J Neurosurg* 2007; 107(5):1004-14.
27. Welter ML, Mallet L, Houeto JL et al. Internal pallidal and thalamic stimulation in patients with Tourette syndrome. *Arch Neurol* 2008; 65(7):952-7.
28. Ackermans L, Duits A, van der Linden C et al. Double-blind clinical trial of thalamic stimulation in patients with Tourette syndrome. *Brain* 2011; 134(Pt 3):832-44.
29. Fontaine D, Lazorthes Y, Mertens P et al. Safety and efficacy of deep brain stimulation in refractory cluster headache: a randomized placebo-controlled double-blind trial followed by a 1-year open extension. *J Headache Pain* 2010; 11(1):23-31.
30. Bussone G, Franzini A, Proietti Cecchini A et al. Deep brain stimulation in craniofacial pain: seven years' experience. *Neurol Sci* 2007; 28 Suppl 2:S146-9.
31. Broggi G, Franzini A, Leone M et al. Update on neurosurgical treatment of chronic trigeminal autonomic cephalgias and atypical facial pain with deep brain stimulation of posterior hypothalamus: results and comments. *Neurol Sci* 2007; 28 Suppl 2:S138-45.
32. Morishita T, Fayad SM, Higuchi MA et al. Deep Brain Stimulation for Treatment-resistant Depression: Systematic Review of Clinical Outcomes. *Neurotherapeutics* 2014.
33. Holtzheimer PE, Kelley ME, Gross RE et al. Subcallosal cingulate deep brain stimulation for treatment-resistant unipolar and bipolar depression. *Arch Gen Psychiatry* 2012; 69(2):150-8.

34. de Koning PP, Figue M, van den Munckhof P et al. Current status of deep brain stimulation for obsessive-compulsive disorder: a clinical review of different targets. *Curr Psychiatry Rep* 2011; 13(4):274-82.
35. Mallet L, Polosan M, Jaafari N et al. Subthalamic nucleus stimulation in severe obsessive-compulsive disorder. *N Engl J Med* 2008; 359(20):2121-34.
36. Denys D, Mantione M, Figue M et al. Deep brain stimulation of the nucleus accumbens for treatment-refractory obsessive-compulsive disorder. *Arch Gen Psychiatry* 2010; 67(10):1061-8.
37. Huff W, Lenartz D, Schormann M et al. Unilateral deep brain stimulation of the nucleus accumbens in patients with treatment-resistant obsessive-compulsive disorder: Outcomes after one year. *Clin Neurol Neurosurg* 2010; 112(2):137-43.
38. Goodman WK, Foote KD, Greenberg BD et al. Deep brain stimulation for intractable obsessive compulsive disorder: pilot study using a blinded, staggered-onset design. *Biol Psychiatry* 2010; 67(6):535-42.
39. Zesiewicz TA, Elble RJ, Louis ED et al. Evidence-based guideline update: treatment of essential tremor: report of the Quality Standards subcommittee of the American Academy of Neurology. *Neurology* 2011; 77(19):1752-5.
40. Zesiewicz TA, Elble R, Louis ED et al. Practice parameter: therapies for essential tremor: report of the Quality Standards Subcommittee of the American Academy of Neurology. *Neurology* 2005; 64(12):2008-20.
41. Pahwa R, Factor SA, Lyons KE et al. Practice Parameter: treatment of Parkinson disease with motor fluctuations and dyskinesia (an evidence-based review): report of the Quality Standards Subcommittee of the American Academy of Neurology. *Neurology* 2006; 66(7):983-95.
42. Zesiewicz TA, Sullivan KL, Arnulf I et al. Practice Parameter: treatment of nonmotor symptoms of Parkinson disease: report of the Quality Standards Subcommittee of the American Academy of Neurology. *Neurology* 2010; 74(11):924-31.
43. Bhidayasiri R, Fahn S, Weiner WJ et al. Evidence-based guideline: treatment of tardive syndromes: report of the Guideline Development Subcommittee of the American Academy of Neurology. *Neurology* 2013; 81(5):463-9.
44. The American Psychiatric Association. 2007 guideline on Treatment of Patients with Obsessive-Compulsive Disorder. 2007. Available online at: [http://www.psychiatryonline.com/pracGuide/pracGuideTopic\\_10.aspx](http://www.psychiatryonline.com/pracGuide/pracGuideTopic_10.aspx)
45. Kennedy SH, Milev R, Giacobbe P et al. Canadian Network for Mood and Anxiety Treatments (CANMAT) Clinical guidelines for the management of major depressive disorder in adults. IV. Neurostimulation therapies. *J Affect Disord* 2009; 117 Suppl 1:S44-53.
46. Muller-Vahl KR, Cath DC, Cavanna AE et al. European clinical guidelines for Tourette syndrome and other tic disorders. Part IV: deep brain stimulation. *Eur Child Adolesc Psychiatry* 2011; 20(4):209-17.
47. National Institute for Health and Care Excellence (NICE). Interventional Procedure Guidance 188. Deep brain stimulation for tremor and dystonia (excluding Parkinson's disease). 2006. Available online at: <http://www.nice.org.uk/nicemedia/pdf/ip/IPG188guidance.pdf>
48. National Institute for Health and Care Excellence (NICE). Interventional Procedure Guidance 382. Deep brain stimulation for refractory chronic pain syndromes (excluding headache) 2011. Available online at: <http://guidance.nice.org.uk/IPG382>
49. National Institute for Health and Care Excellence (NICE). Interventional Procedure Guidance 381. Deep brain stimulation for intractable trigeminal autonomic cephalgias. 2011. Available online at: <http://www.nice.org.uk/IPG381>
50. National Institute for Health and Care Excellence (NICE). Interventional Procedure Guidance 416. Deep brain stimulation for refractory epilepsy. 2012. Available online at: <http://guidance.nice.org.uk/IPG416>
51. National Institute for Health and Care Excellence (NICE). Interventional Procedure Guidance 19. Deep brain stimulation for Parkinson's disease. 2003. Available online at: <http://www.nice.org.uk/nicemedia/pdf/ip/IPG019guidance.pdf>
52. National Institute for Health and Care Excellence (NICE). Parkinson's disease: Diagnosis and management in primary and secondary care. Clinical Guideline 35. 2006. Available online at: <http://www.nice.org.uk/nicemedia/live/10984/30088/30088.pdf>

53. Centers for Medicare and Medicaid. NCD for Deep Brain Stimulation for Essential Tremor and Parkinson's Disease (160.24). 2003. Available online at: <http://www.cms.gov/medicare-coverage-database/search/document-id-search-results.aspx?DocID=160.24&bc=qAAAAAAAAAA&>

### **Billing Coding/Physician Documentation Information**

---

<b>61860</b>	Craniectomy or craniotomy for implantation of neurostimulator electrodes, cerebral, cortical
<b>61863</b>	Twist drill, burr hole, craniotomy, or craniectomy with stereotactic implantation of neurostimulator electrode array in subcortical site (eg, thalamus, globus pallidus, subthalamic nucleus, periventricular, periaqueductal gray), without use of intraoperative microelectrode recording; first array
<b>61864</b>	Twist drill, burr hole, craniotomy, or craniectomy with stereotactic implantation of neurostimulator electrode array in subcortical site (eg, thalamus, globus pallidus, subthalamic nucleus, periventricular, periaqueductal gray), without use of intraoperative microelectrode recording; each additional array (List separately in addition to primary procedure)
<b>61867</b>	Twist drill, burr hole, craniotomy, or craniectomy with stereotactic implantation of neurostimulator electrode array in subcortical site (eg, thalamus, globus pallidus, subthalamic nucleus, periventricular, periaqueductal gray), with use of intraoperative microelectrode recording; first array
<b>61868</b>	Twist drill, burr hole, craniotomy, or craniectomy with stereotactic implantation of neurostimulator electrode array in subcortical site (eg, thalamus, globus pallidus, subthalamic nucleus, periventricular, periaqueductal gray), with use of intraoperative microelectrode recording; each additional array (List separately in addition to primary procedure)
<b>61875</b>	Craniectomy for implantation of neurostimulator electrodes, cerebellar; subcortical
<b>61880</b>	Revision or removal of intracranial neurostimulator electrodes
<b>61885</b>	Incision and subcutaneous placement of cranial neurostimulator pulse generator or receiver, direct or inductive coupling; with connection to a single electrode array.
<b>61886</b>	Incision and subcutaneous placement of cranial neurostimulator pulse generator or receiver, direct or inductive coupling; with connection to two or more electrode arrays
<b>61888</b>	Revision or removal of cranial neurostimulator pulse generator or receiver
<b>95961</b>	Functional cortical and subcortical mapping by stimulation and/or recording of electrodes on brain surface, or of depth electrodes, to provoke seizures or identify vital brain structures; initial hour of physician attendance
<b>95962</b>	as above; each additional hour
<b>95970</b>	Electronic analysis of implanted neurostimulator pulse generator system, brain, spinal cord or peripheral pulse generator / transmitter, without reprogramming
<b>95971</b>	as above, with intraoperative or subsequent programming
<b>95972</b>	as above, first hour of programming
<b>95973</b>	as above, each additional 30 minutes
<b>95974</b>	as 95970- , with or without nerve interface testing, first hour
<b>95975</b>	as 95974 – each additional 30 minutes
<b>95978</b>	Electronic analysis of implanted neurostimulator pulse generator system (eg, rate, pulse amplitude and duration, battery status, electrode selectability and polarity, impedance and patient compliance measurements), complex deep brain neurostimulator pulse generator/transmitter, with initial or subsequent programming; first hour
<b>95979</b>	Electronic analysis of implanted neurostimulator pulse generator system (eg, rate, pulse amplitude and duration, battery status, electrode selectability and polarity, impedance and patient compliance measurements), complex deep brain neurostimulator pulse generator/transmitter, with initial or subsequent programming; each additional 30 minutes after first hour (List separately in addition to code for primary procedure)
<b>L8679</b>	Implantable neurostimulator, pulse generator, any type
<b>L8680</b>	Implantable neurostimulator electrode, each
<b>L8682</b>	Implantable neurostimulator radiofrequency receiver
<b>L8683</b>	Radiofrequency transmitter (external) for use with implantable neurostimulator

	radiofrequency receiver
<b>L8685</b>	Implantable neurostimulator pulse generator, single array, rechargeable, includes extension
<b>L8686</b>	Implantable neurostimulator pulse generator, single array, non-rechargeable, includes extension
<b>L8687</b>	Implantable neurostimulator pulse generator, dual array, rechargeable, includes extension
<b>L8688</b>	Implantable neurostimulator pulse generator, dual array, non-rechargeable, includes extension
<b>C1787</b>	Patient programmer, neurostimulator
<b>C1820</b>	Generator, neurostimulator (implantable), with rechargeable battery and charging system
<b>C1897</b>	Lead, neurostimulator test kit (implantable)

The coding for deep brain stimulation consists of a series of CPT codes describing the various steps of the procedure; i.e., implantation of the electrodes, implantation of the pulse generator, intraoperative monitoring and programming of the electrodes, postoperative neuroprogramming.

## **Additional Policy Key Words**

N/A

## **Policy Implementation/Update Information**

8/1/01	New policy.
8/1/02	Policy statement revised to indicate bilateral stimulation may be considered medically necessary with certain criteria.
8/1/03	No policy statement changes.
8/1/04	Policy statement revised to include the following as medically necessary: Patients aged greater than 7 years with chronic, intractable (drug refractory) primary dystonia, including generalized and/or segmental dystonia, hemidystonia and cervical dystonia (torticollis) in patients seven years of age or above
8/1/05	Policy statement revised to include the following as investigational: For the treatment of chronic cluster headaches.
3/1/06	Interim change, added CPT codes
8/1/06	No policy statement changes.
8/1/07	No policy statement changes.
8/1/08	Policy statement revised to include deep brain stimulation for the treatment of other psychiatric or neurologic disorders, including but not limited to Tourette syndrome, depression, obsessive compulsive disorder and epilepsy as investigational.
8/1/09	No policy statement changes.
8/1/10	No policy statement changes.
8/1/11	No policy statement changes.
8/1/12	No policy statement changes.
8/1/13	Anorexia nervosa, alcohol addiction and chronic pain added as investigational.
8/1/14	Added Alzheimer disease to Investigational Statement.

State and Federal mandates and health plan contract language, including specific provisions/exclusions, take precedence over Medical Policy and must be considered first in determining eligibility for coverage. The medical policies contained herein are for informational purposes. The medical policies do not constitute medical advice or medical care. Treating health care providers are independent contractors and are neither employees nor agents Blue KC and are solely responsible for diagnosis, treatment and medical advice. No part of this publication may be reproduced, stored in a retrieval system or transmitted, in any form or by any means, electronic, photocopying, or otherwise, without permission from Blue KC.