

Medical Policy



Title: Treatment of Hyperhidrosis

Professional

Original Effective Date: September 23, 2009
Revision Date(s): January 26, 2010;
March 31, 2014
Current Effective Date: March 31, 2014

Institutional

Original Effective Date: September 23, 2009
Revision Date(s): January 26, 2010
March 31, 2014
Current Effective Date: March 31, 2014

State and Federal mandates and health plan member contract language, including specific provisions/exclusions, take precedence over Medical Policy and must be considered first in determining eligibility for coverage. To verify a member's benefits, contact [Blue Cross and Blue Shield of Kansas Customer Service](#).

The BCBSKS Medical Policies contained herein are for informational purposes and apply only to members who have health insurance through BCBSKS or who are covered by a self-insured group plan administered by BCBSKS. Medical Policy for FEP members is subject to FEP medical policy which may differ from BCBSKS Medical Policy.

The medical policies do not constitute medical advice or medical care. Treating health care providers are independent contractors and are neither employees nor agents of Blue Cross and Blue Shield of Kansas and are solely responsible for diagnosis, treatment and medical advice.

If your patient is covered under a different Blue Cross and Blue Shield plan, please refer to the Medical Policies of that plan.

DESCRIPTION

Hyperhidrosis, or excessive sweating, can lead to impairments in psychological and social functioning. Various treatments for hyperhidrosis are available, such as topical agents, oral medications, botulinum toxin, and surgical procedures.

Background

Hyperhidrosis may be defined as excessive sweating, beyond a level required to maintain normal body temperature in response to heat exposure or exercise. It can be classified as either primary or secondary. Primary focal hyperhidrosis is idiopathic in nature, typically involving the hands (palmar), feet (plantar), or axillae (underarms). Secondary hyperhidrosis can result from a variety of drugs, such as tricyclic antidepressants,

selective serotonin reuptake inhibitors (SSRIs), or underlying diseases/conditions, such as febrile diseases, diabetes mellitus, or menopause.

Secondary hyperhidrosis is usually generalized or craniofacial sweating. Secondary gustatory hyperhidrosis is excessive sweating on ingesting highly spiced foods. This trigeminovascular reflex typically occurs symmetrically on scalp or face and predominately over forehead, lips, and nose. Secondary facial gustatory sweating, in contrast, is usually asymmetrical and occurs independently of the nature of the ingested food. This phenomenon frequently occurs after injury or surgery in the region of the parotid gland. Frey's syndrome is an uncommon type of secondary gustatory hyperhidrosis that arises from injury to or surgery near the parotid gland resulting in damage to the secretory parasympathetic fibers of the facial nerve. After injury, these fibers regenerate, and miscommunication occurs between them and the severed postganglionic sympathetic fibers that supply the cutaneous sweat glands and blood vessels. The aberrant connection results in gustatory sweating and facial flushing with mastication. Aberrant secondary gustatory sweating follows up to 73% of surgical sympathectomies and is particularly common after bilateral procedures.

The consequences of hyperhidrosis are primarily psychosocial in nature. Symptoms such as fever, night sweats, or weight loss require further investigation to rule out secondary causes. Sweat production can be assessed with the minor starch iodine test, which is a simple qualitative measure to identify specific sites of involvement.

A variety of therapies have been investigated for primary hyperhidrosis, including topical therapy with aluminum chloride, oral anticholinergic medications, iontophoresis, intradermal injections of botulinum toxin, endoscopic transthoracic sympathectomy, and surgical excision of axillary sweat glands. Treatment of secondary hyperhidrosis focuses on treatment of the underlying cause, such as discontinuing certain drugs or hormone replacement therapy as a treatment of menopausal symptoms.

Botulinum toxin is a potent neurotoxin that blocks cholinergic nerve terminals; symptoms of botulism include cessation of sweating. Therefore, intracutaneous injections have been investigated as a treatment of gustatory hyperhidrosis and focal primary hyperhidrosis, most frequently involving the axillae or palms. The drawback of this approach is the need for repeated injections, which have led some to consider surgical approaches.

Surgical treatment options include removal of the eccrine glands and/or interruption of the sympathetic nerves. Eccrine sweat glands produce an aqueous secretion, the overproduction of which is primarily responsible for hyperhidrosis. These glands are innervated by the sympathetic nervous system. Surgical removal has been performed in patients with severe isolated axillary hyperhidrosis.

Various surgical techniques of sympathectomy may also be tried. The second (T2) and third (T3) thoracic ganglia are responsible for palmar hyperhidrosis, the fourth (T4)

thoracic ganglion controls axillary hyperhidrosis, and the first (T1) thoracic ganglion controls facial hyperhidrosis. Thoracic sympathectomy has been investigated as a potentially curative procedure, primarily for combined palmar and axillary hyperhidrosis that is unresponsive to non-surgical treatments. While accepted as an effective treatment, sympathectomy is not without complications. In addition to the immediate surgical complications of pneumothorax or temporary Horner's syndrome, compensatory sweating on the trunk generally occurs in a majority of patients, with different degrees of severity. Medical researchers have investigated whether certain approaches, e.g., T3 versus T4 sympathectomy, result in less compensatory sweating, but there remains a lack of consensus about which approach best minimizes the risk of this side effect. In addition, with lumbar sympathectomy for plantar hyperhidrosis, there has been concern about the risk of post-operative sexual dysfunction in men and women.

The outcome of different surgical and medical treatment modalities is best assessed by using a combination of tools. Quantitative tools include gravimetry, evaporimetry, and Minor's starch iodine test. Qualitative assessment tools include general health surveys and hyperhidrosis-specific surveys. Of these, the Hyperhidrosis Disease Severity Scale (HDSS) has been found to have a good correlation to other assessment tools and to be practical in the clinical setting.

Regulatory Status

Drysol™ (aluminum chloride [hexahydrate] 20% topical solution, Person and Covey, Inc.) is approved by the U.S. Food and Drug Administration (FDA) to be used as an aid in the management of hyperhidrosis (axillae, palmar, plantar, and craniofacial); it is available by prescription.

In 2004 the FDA approved botulinum toxin type A (Botox®) to treat primary axillary hyperhidrosis (severe underarm sweating) that cannot be managed by topical agents. In 2009, this product was renamed to OnabotulinumtoxinA. Other FDA-approved botulinum toxin products include:

- 2000: RimabotulinumtoxinB, marketed as Myobloc® (Solstice Neurosciences)
- 2009: AbobotulinumtoxinA, marketed as Dysport® (Medicis Pharmaceutical Corporation, Scottsdale, AZ)
- 2010: IncobotulinumtoxinA, marketed as Xeomin® (Merz Pharmaceuticals)

None of these other botulinum toxin products are indicated for treatment of hyperhidrosis.

On July 31, 2009, the FDA approved the following revisions to the prescribing information of botulinum toxin products:

- "A *Boxed Warning* highlighting the possibility of experiencing potentially life-threatening distant spread of toxin effect from injection site after local injection.
- A Risk Evaluation and Mitigation Strategy (REMS) that includes a *Medication Guide* to help patients understand the risk and benefits of botulinum toxin products.

- Changes to the established drug names to reinforce individual potencies and prevent medication errors. The potency units are specific to each botulinum toxin product, and the doses or units of biological activity cannot be compared or converted from one product to any other botulinum toxin product. The new established names reinforce these differences and the lack of interchangeability among products.”

In January 2011, the miraDry® System (Miramar Labs, Inc.; Sunnydale, CA) was cleared by the FDA through the 510(k) process for treating primary axillary hyperhidrosis. This is a microwave device designed to heat tissue at the dermal-hypodermal interface, the location of the sweat glands. Treatment consists of 2 sessions of approximately one hour in duration. Sessions occur in a physician’s office and local anesthetic is used

POLICY

A. Primary Focal Hyperhidrosis

Treatment of primary hyperhidrosis (see definition of primary hyperhidrosis in Policy Guidelines) may be considered **medically necessary** with any of the following medical complications:

1. acrocyanosis of the hands
or
2. history of recurrent skin maceration with bacterial or fungal infections
or
3. history of recurrent secondary infections
or
4. history of persistent eczematous dermatitis in spite of medical treatments with topical dermatological or systemic anticholinergic agents

Treatments for Primary Focal Hyperhidrosis

Focal Regions	Treatments Considered Medically Necessary	Treatments Considered Investigational
Axillary	<ul style="list-style-type: none"> ▪ aluminum chloride 20% solution* ▪ botulinum toxin* for severe primary axillary hyperhidrosis that is inadequately managed with topical agents*, in patients 18 years and older ▪ endoscopic transthoracic sympathectomy (ETS) and surgical excision of axillary sweat glands, if conservative treatment (i.e., aluminum chloride or botulinum toxin, individually and in combination) has failed 	<ul style="list-style-type: none"> ▪ axillary liposuction ▪ iontophoresis ▪ microwave treatment

Focal Regions	Treatments Considered Medically Necessary	Treatments Considered Investigational
Palmar	<ul style="list-style-type: none"> ▪ aluminum chloride 20% solution* ▪ botulinum toxin A products for severe primary palmar hyperhidrosis that is inadequately managed with topical agents, in patients 18 years and older ▪ endoscopic transthoracic sympathectomy (ETS), if conservative treatment (i.e., aluminum chloride or botulinum toxin type A, individually and in combination) has failed 	<ul style="list-style-type: none"> ▪ RimabotulinumtoxinB ▪ iontophoresis ▪ microwave treatment ▪ radiofrequency ablation
Plantar	<ul style="list-style-type: none"> ▪ aluminum chloride 20% solution* 	<ul style="list-style-type: none"> ▪ botulinum toxin ▪ iontophoresis ▪ lumbar sympathectomy ▪ microwave treatment
Craniofacial	<ul style="list-style-type: none"> ▪ aluminum chloride 20% solution* ▪ endoscopic transthoracic sympathectomy (ETS), if conservative treatment (i.e., aluminum chloride) has failed 	<ul style="list-style-type: none"> ▪ botulinum toxin ▪ iontophoresis ▪ microwave treatment

*FDA approved indication. Note: In the case of botulinum toxin, FDA approved for at least one botulinum toxin product.

B. Secondary Hyperhidrosis

Secondary Gustatory Hyperhidrosis

1. The following treatments would be considered **medically necessary** for the treatment of severe gustatory hyperhidrosis:
 - a. aluminum chloride 20% solution*
 - b. surgical options (i.e., tympanic neurectomy), if conservative treatment has failed.

*FDA approved indication. Note: In the case of botulinum toxin, FDA approved for at least one botulinum toxin product.

2. The following treatments are considered **experimental / investigational** as a treatment for severe gustatory hyperhidrosis including, but not limited to:
 - a. botulinum toxin is considered investigational for treatment of gustatory hyperhidrosis
 - b. iontophoresis
3. Treatment of hyperhidrosis is considered **not medically necessary** in the absence of functional impairment or medical complications.

Policy Guidelines

A multispecialty working group defines primary focal hyperhidrosis as a condition that is characterized by visible, excessive sweating of at least 6 months' duration without apparent cause and with at least 2 of the following features:

- a. bilateral and relatively symmetric sweating,
- b. impairment of daily activities,
- c. frequency of at least once per week,
- d. age at onset younger than 25 years,
- e. positive family history, and
- f. cessation of focal sweating during sleep. (1)

RATIONALE

This policy was updated with searches of the MEDLINE database. The most recent literature search was performed for the period February 2012 through March 18, 2013. Following is a summary of the key literature to date:

Iontophoresis

The published literature regarding iontophoresis as a treatment of hyperhidrosis is sparse. A 2003 TEC Assessment on iontophoresis concluded that the evidence was insufficient to determine whether the effects of iontophoresis for the treatment of hyperhidrosis exceed those of placebo or an alternative treatment. (3) The TEC Assessment investigators identified only 3 small studies (n=10, 11, and 18, respectively), all of which were conducted in patients with palmar hyperhidrosis.

Since the 2003 TEC Assessment, no randomized controlled trials (RCTs) evaluating tap water iontophoresis for treating hyperhidrosis have been published. Subsequent case series include a 2013 study from Ireland with 28 patients. (4) Patients received a minimum of 9 treatments over 21 days in a clinical setting. Twenty of the 25 patients (80%) for whom data were available after hospital administration of tap water iontophoresis reported a moderate or great amount of improvement in symptoms and a moderate or great improvement in quality of life (according to a Disease Life Quality Index). After the initial course of treatment, 13 patients chose to purchase a home iontophoresis device. Eight of the 13 (62%) reported that home iontophoresis was 'much less effective' than hospital treatment. No other outcomes e.g., symptom improvement, were reported following home use.

Conclusion: There is insufficient evidence, consisting of only small case series, that iontophoresis is an effective treatment of hyperhidrosis. Controlled trials are needed to determine whether iontophoresis improves the net health outcome in patients with hyperhidrosis.

Botulinum toxin type A

A considerable body of published literature addresses botulinum toxin injection of the treatment of axillary and palmar hyperhidrosis and substantiates the efficacy of this treatment. (5-13) Studies include multiple randomized placebo-controlled trials evaluating Botox, a botulinum toxin type A product. In addition, another botulinum toxin A product, Dysport, has been evaluated in RCTs for treatment of axillary hyperhidrosis (13) and palmar hyperhidrosis. (8) Moreover, a small

RCT published in 2007 compared Botox and Dysport and found similar levels of efficacy and safety with the two products. (10)

One of the larger RCTs was published in 2007. (12) This study was an industry-sponsored multicenter double-blind, placebo-controlled efficacy and safety study of botulinum toxin type A in patients with persistent bilateral primary axillary hyperhidrosis. Enrollment criteria included a resting sweat production of at least 50 mg/axilla in 5 minutes and a rating of 3 or 4 (underarm sweating barely tolerable or intolerable, and frequently or always interferes with daily activities) on the Hyperhidrosis Disease Severity Scale (HDSS). A total of 322 patients were randomized to receive 50 U, or 75 U of Botox or placebo. Retreatment after 4 weeks was allowed in subjects with at least 50 mg of sweat (per axilla) over 5 minutes and an HDSS score of 3 or 4. Following the first injection, 75% of subjects in the Botox groups showed at least a 2-point improvement in the HDSS, compared with 25% of subjects in the placebo group. Sweat production decreased by 87% (75 U), 82% (50 U), and 33% (vehicle). (Similar results were obtained in patients requiring a second treatment.) The median duration of effect was 197, 205, and 96 days (75 U, 50 U, and vehicle, respectively). Seventy-eight percent of subjects (252) completed the 52-week study; 96 of 110 (87%) in the 75-U group, 83 of 104 (80%) in the 50-U group, and 73 of 108 (68%) in the control group. Intent-to-treat analysis at 52 weeks showed a responder rate (greater than 2-point improvement on the HDSS) for 54 (49%) subjects in the 75-U group, 57 (55%) in the 50-U group, and 6 (6%) in the placebo group. Injection-site pain was reported in about 10% of all groups, with a mean duration of 2.4 days (10-day maximum).

No placebo-controlled RCTs were identified evaluating the safety and efficacy of the newest formulation of botulinum toxin A, Xeomin. In 2010, Dressler published a double-blind RCT from Germany comparing Xeomin to Botox for treating primary axillary hyperhidrosis. (14) Forty-six patients with bilateral axillary hyperhidrosis and a previously stable Botox treatment for at least 2 years received 50 MU of Botox in one axilla and 50 MU Xeomin in the other axilla. All patients completed the study. A total of 41 of 46 (89%) patients reported the therapeutic effect as excellent and 5 (11%) as good. The mean reported duration of therapeutic effect was 3.2 months. According to patient self-report in structured interviews, there were no side-to-side differences in therapeutic effect including onset latency, extent and duration and no differences in injection site pain. Moreover, clinical examination did not identify any side-to-side differences in the diffuse sweating pattern.

There is less evidence in support of botulinum toxin type A for treating plantar hyperhidrosis. No RCTs or large uncontrolled studies were identified; most published studies are case reports or small case series.

The evidence evaluating botulinum toxin A use for gustatory hyperhidrosis as a result of Frey's syndrome includes noncontrolled or nonrandomized studies, all showing favorable treatment outcomes. The patient inclusion criteria were variable across the studies and case reports; ages varied (16 to 87 years); patients had undergone varied types of parotid surgery (i.e., bilateral, partial); not all studies documented gustatory sweating with Minor's starch test as part of the patient screening.

Conclusions: Multiple RCTs support the efficacy and safety of botulinum toxin A for treating severe axillary and palmar hyperhidrosis. There is a lack of RCTs on use of botulinum toxin A for plantar hyperhidrosis and gustatory hyperhidrosis.

Botulinum toxin type B

There was one placebo-controlled randomized trial on botulinum toxin B (Myobloc) for treating primary axillary hyperhidrosis and one on palmar hyperhidrosis. Both studies were by Baumann and colleagues and were published in 2005; neither discussed whether patients had failed previous treatments for hyperhidrosis. The study on axillary hyperhidrosis included 20 participants; they received subcutaneous injections of Myobloc (2,500 U or 0.5 mL per axilla) (n=15) or placebo (n=5). (15) Patients who received placebo were offered Myobloc at subsequent injections. One patient in the placebo group did not return for follow-up and another responded to placebo and did not return for a subsequent Myobloc injection. Data were available on Myobloc efficacy for the remaining 18 participants (15 in the initial Myobloc group and 3 crossovers). There was a statistically significant improvement in axillary hyperhidrosis according to patient and physician subjective assessment from baseline (before receiving an active injection) to Day 30. Details on the efficacy outcomes were not reported. The mean length of time to return to baseline levels of sweating in these 18 patients was 151 days (range 66 to 243 days). Sixteen participants reported 61 adverse events over the course of the study. Five of 61 adverse events (8.2%) were determined to be definitely related to the study; 4 axillary bruising events and 1 instance of pain at the injection site. Eleven adverse events (18%) were determined to be probably related to study treatment; dry eyes (n=3), dry mouth (n=5) and indigestion (n=3). Flu-like symptoms were reported by 6 of 20 patients (30%); however, the study period coincided with flu season. Note that the authors did not compare the active treatment and placebo groups in their analysis.

The RCT on Myobloc for treatment of palmar hyperhidrosis included 20 participants with excessive palmar sweating. Fifteen participants received injections of Myobloc (50,000 U per palm) and 5 received placebo. (16) Nonresponders were offered an injection of Myobloc at day 30. At day 30, the two quality-of-life measures were significantly higher in the Myobloc group compared to the control group. However, there was not a statistically significant difference in efficacy in the physician analysis of the palmar iodine starch test at day 30 ($p=0.56$). No further details were provided on the efficacy outcome measures described above. The mean duration of action according to self-report in 17 patients (15 in the initial treatment group and 2 who crossed-over from the placebo group) was 3.8 months (range, 2.3 to 4.9 months). Participants were asked about specific adverse events. Eighteen of 20 (90%) reported dry mouth/throat, 12 (60%) reported indigestion, 12 (60%) reported excessively dry hands, 12 (60%) reported muscle weakness, and 10 (50%) reported decreased grip strength. Both studies by Baumann and colleagues were limited by a small sample sizes and limited or no comparative data.

A small randomized trial by Frasson and colleagues in Italy that compared botulinum toxin type A and type B for treating axillary hyperhidrosis was published in 2011. (17) This study included 10 patients with idiopathic focal axillary hyperhidrosis that was unresponsive to other non-surgical treatments. Patients received 50 U botulinum toxin A in one axilla and 2,500 U botulinum toxin B in the contralateral axilla. Gravimetry was performed at baseline and follow-up as an objective measurement of sweat production. In addition, the sweat area was photographed. At each follow-up point, the decrease in sweat weight from baseline was significantly greater on the botulinum toxin B side compared to the botulinum toxin A side. For example, after 1 month, the sweat weight in 5 minutes was 13% of the baseline value on the botulinum toxin A side and 4% of the baseline value on the botulinum toxin B side ($p=0.049$). By 6 months, the sweat weight returned to 91% of baseline on the botulinum toxin A side and 56% of baseline weight on the botulinum toxin B side ($p=0.02$). Findings were similar for sweating area. All patients tolerated injections of

botulinum toxin types A and B well and none reported systemic adverse effects. The authors commented that this study used a higher dosage of botulinum toxin B than previous studies.

Conclusions: There are few RCTs evaluating botulinum toxin type B for treating hyperhidrosis. One small RCT did not clearly demonstrate the efficacy of botulinum toxin type B in patients with palmar hyperhidrosis. Two RCTs support the efficacy of this treatment for patients with axillary hyperhidrosis.

Microwave treatment

A 2012 RCT evaluated a microwave device for treating hyperhidrosis. (18) This device applies microwave energy to superficial skin structures with the intent of inducing thermolysis of the eccrine and apocrine sweat glands. This industry-sponsored double-blind study randomized 120 adults with primary axillary hyperhidrosis in a 2-to-1 ratio to active (n=81) or sham (n=39) treatment. Treatment consisted of 2 sessions, separated by approximately 2 weeks. Patients who responded adequately after 1 session or declined further treatment did not need to undergo the second session, and a third procedure was allowed within 30 days for participants who still had a high level of sweating after 2 sessions. All patients in the sham group had 2 sessions. In the active treatment group, 11 individuals (9%) had only 1 session and 10 (8%) had a third procedure. The primary efficacy endpoint was a score of 1 (underarm sweating never noticeable) or 2 (underarm sweating tolerable) on the Hyperhidrosis Disease Severity Scale at the 30-day follow-up; HDSS score at 6 months was a secondary outcome. A total of 101/120 (84%) completed the study. At 30 days, 89% of the active treatment group and 54% of the sham group had an HDSS score of 1 or 2; $p < 0.001$. At 6 months, 67% of the active treatment group and 44% of the sham group had an HDSS score of 1 or 2; the difference between groups remained statistically significant, $p = 0.02$. Unblinding occurred at 6 months. Twelve-month data were available for the active treatment group only; 69% reported an HDSS score of 1 or 2. There were 45 procedure-related adverse events in 23 (28%) of the active treatment group and 5 (13%) of the sham group. The most frequently reported adverse event was altered sensation; no serious adverse events were reported. Compensatory sweating was reported by 2 individuals in each group and had a mean duration of 52 days. The authors noted that study data provided an opportunity to identify areas for improvement of the treatment protocol including waiting longer between treatments and using a higher dose of energy at the second session.

A 2012 industry-sponsored case series reported on 31 patients with primary axillary hyperhidrosis who were treated with microwave therapy using the miraDry system. (19) All patients had an HDSS score of 3 or 4 at baseline. The primary efficacy outcome, the proportion of patients whose HDSS decreased to 1 or 2 was 28 (90%) at 6 months and 12 months after treatment. Longer-term skin-related adverse effects (that all resolved over time) were altered sensation in the skin of the axillae (65% of patients, median duration, 37 days) and palpable bumps under the skin of the axillae (71% of patients, median duration, 41 days).

Conclusions: One RCT and case series provide insufficient evidence that microwave treatment improves the health outcome for primary focal hyperhidrosis. The RCT reported short-term benefit of microwave treatment in reducing hyperhidrosis, but also reports a high rate of skin-related side effects such as pain and altered sensation. Additional controlled studies with long-term follow-up in the treatment and control groups, a longer period of blinding, and a consistent treatment protocol are needed to confirm the efficacy of this treatment and to better define the risk/benefit ratio.

Radiofrequency ablation

A 2013 study evaluated radiofrequency ablation as a treatment option for patients with severe bilateral palmar hyperhidrosis resistant to conservative treatment. (20) The study was conducted in Turkey and retrospectively reviewed outcomes after radiofrequency ablation (n=48) or transthoracic sympathectomy (n=46). Patients were not randomized to treatment group. After the mean of 15-month follow-up, palmar hydrosis was absent in 36 patients (75%) in the radiofrequency ablation group and 44 patients (96%) in the surgery group. The difference in outcomes was statistically significant between groups, favoring the surgical intervention ($p<0.01$). Six patients in each group reported moderate or severe compensatory sweating ($p=0.78$).

Conclusions: One nonrandomized comparative study represents insufficient evidence on radiofrequency ablation as a treatment of hyperhidrosis. In this single available study, radiofrequency ablation was found to be inferior to surgical sympathectomy.

Surgical interventions

Tympanic neurectomy for gustatory hyperhidrosis

Review articles by Clayman et al. (21) and de Bree et al. (22) describe the various medical and surgical treatments for Frey's syndrome. Tympanic neurectomy is described as a treatment for Frey's syndrome, with satisfactory control reported in 82% of patients. In addition, this surgical treatment is generally definitive without a need for repeated interventions.

Sweat gland excision for primary focal hyperhidrosis

Surgery may involve removal of the subcutaneous sweat glands without removal of any skin, limited excision of skin, and removal of surrounding subcutaneous sweat glands, or a more radical excision of skin and subcutaneous tissue en bloc. (23) Depending on the completeness of surgical excision, the treatment is effective in 50–95% of patients.

Transthoracic sympathectomy for primary focal hyperhidrosis

Several RCTs and one meta-analysis have compared different approaches to surgery; there were no sham-controlled RCTs. In 2011, Deng and colleagues published a meta-analysis of data from RCTs and observational studies published to 2010 evaluating thoracoscopic sympathectomy for patients with palmar hyperhidrosis. (24) The authors pooled outcome data from different approaches to sympathectomy, i.e., single-ganglia blockage (T2, T3, or T4), and multi-ganglia blockage (T2-3, T2-4, or T3-4). (Note: T refers to rib). Based on these analyses, they concluded that T3 (11 studies) and T3-4 (2 studies) had the “best” clinical efficacy i.e., postoperative resolution of symptoms. The T3 approach resulted in a 97.9% pooled efficacy rate, and the T3-4 approach resulted in a 100% pooled efficacy rate. In the studies for which data were available, the pooled rate of postoperative compensatory sweating was 40% after T3 surgery. Data on compensatory sweating after T3-4 surgery was only available from one study with 60 patients; a pooled analysis could not be performed.

Subsequent RCTs have also compared levels of sympathectomy. For example, a 2011 study by Baumgartner and colleagues included 121 patients with disabling palmoplantar hyperhidrosis. (25) Patients were randomized to receive bilateral sympathectomy over T2 (n=61 patients) or T3 (n=60 patients). Six of 121 (5%) patients, 3 in each group, were considered treatment failures i.e., had recurrent palmar sweating to a bothersome level. There were no significant differences between groups in the reported subjective change in plantar or axillary sweating after surgery. At 6 months, the mean level of compensatory sweating (0 to 10 severity scale) was 4.7 (standard

deviation [SD]=2.7) for the T2 group and 3.8 (SD=2.8) for the T3 group (p =not significant). Similarly, at 1 year, the mean severity rating of compensatory sweating was 4.7 (SD=2.5) in the T2 group and 3.7 (SD=2.8) in the T3 group; $p=0.09$. Another study was published by Ishy and colleagues in Brazil in which surgery at the T3 and T4 levels was compared. (26) This study included 20 patients with palmar hyperhidrosis. All patients experienced complete bilateral remission of palmar sweating after 1 year of follow-up. The level of compensatory sweating did not differ significantly between groups at 1 week, 1 month, or 6 months, but at 1 year, there was a significantly higher rate in the T3 compared to the T4 group (20/20, 100% in the T3 group and 15/20, 75% in the T4 group, $p=0.47$)

There is also a large amount of data from case series on transthoracic sympathectomy for treating primary focal hyperhidrosis. (27-32) Case series generally report high success rates for palmar and axillary hyperhidrosis, although there are potential adverse effects, most commonly compensatory sweating.

For example, in 2010, Wait and colleagues published a retrospective analysis of prospectively collected data on patients who underwent bilateral thoracoscopic sympathectomy for hyperhidrosis. (31) Data were available on 322 of 348 (93%) of patients who underwent surgery. Patients' previous use of nonsurgical hyperhidrosis treatments was not reported. Complete resolution of symptoms was reported by 300 of 301 (99.7%) with palmar hyperhidrosis, 136 of 186 (73%) with axillary hyperhidrosis, 27 of 30 (90%) with craniofacial hyperhidrosis, and 19 of 197 (9.6%) with plantar hyperhidrosis. Compensatory sweating was reported by a total of 201 of 322 (62%) patients. The compensatory sweating was severe in 20 (6.2%) of patients and mild or moderate in 181 (56.2%) of patients. It is worth noting that thoracoscopic sympathectomy was performed in some cases of plantar hyperhidrosis and that there was a low rate of success. In addition, when reporting rates of compensatory sweating, the authors did not distinguish between mild and moderate levels of symptoms, although these could have different clinical implications for the patient.

In 2011, Smidfelt and Drott in Sweden reported on long-term outcomes after transthoracic sympathectomy. (32) Of 3,015 patients who had been treated with endoscopic thoracic sympathectomy for hyperhidrosis and/or facial blushing, 1,700 (56%) responded to a written survey after a mean of 14.6 (SD=2.4) years. A total of 85.1% of respondents reported that they had a satisfactory and lasting effect of the surgery. Sweating and/or blushing recurred and was considered a problem in 8.1%, and 6.9% reported no initial effect or a poor effect. Compensatory sweating was considered troublesome by 299 (17.6%), annoying by 409 (24.1%), severe by 367 (21.6%), and incapacitating by 190 (11.2%). Nearly half of the patients who underwent surgery did not respond to the survey; their outcomes may have been different from those of study respondents.

A 2013 series reported on complications after thoracic sympathectomy in 1,731 patients with palmar, axillary or craniofacial hyperhidrosis. (33) Thirty days after surgery, 1,531 (88.4%) of patients reported compensatory sweating. Among the 1,531 patients, compensatory sweating was mild in 473 (31%), moderate in 642 (42%) and severe in 416 (27%). Gustatory sweating was reported by 334 of the 1,731 (19%) patients.

Conclusions: RCTs and a meta-analysis of RCTs support the efficacy of transthoracic sympathectomy at various levels for palmar and axillary hyperhidrosis. These data are

complemented by case series which have found high efficacy rates, but also high rates of compensatory sweating for these conditions. There is insufficient evidence in support of transthoracic sympathectomy for treating plantar hyperhidrosis; case series found lower rates of efficacy for plantar compared to axillary or palmar hyperhidrosis, and there are concerns for side effects in sexual functioning.

Endoscopic lumbar sympathectomy for primary plantar hyperhidrosis

No RCTs were identified but several case series were identified. A 2009 series by Rieger and colleagues from Austria evaluated surgery results in 90 patients (59 men, 31 women with severe plantar hyperhidrosis). (34) Thirty-seven patients (41%) had only plantar hyperhidrosis, and 53 (59%) had plantar and palmar hyperhidrosis. All patients had previously used other treatments including topical aluminum chloride therapy. There were a total of 178 procedures—90 on the right-side and 88 on the left side. The technique involved resecting a segment of the sympathetic trunk between the third and fourth lumbar bodies together with the ganglia (L3 and/or L4). After a mean follow-up of 24 months (range: 3 to 45), hyperhidrosis was eliminated in 87 of 90 patients (97%). Postoperative neuralgia occurred in 38 (42%) patients between the seventh and eighth day. The pain lasted less than 4 weeks in 11 patients, 1-3 months in 19 patients, 4-12 months in 5 patients, and more than 12 months in 3 patients. Three men reported temporary sexual symptoms; one was incapable of ejaculation for 2 months. None of the women reported postoperative sexual dysfunction.

In 2010, Reisfeld reported on results of a U.S.-based study from a specialized hyperhidrosis clinic in which bilateral endoscopic lumbar sympathectomy was performed in 63 patients with focal plantar hyperhidrosis. (35) There were 13 (21%) male patients and 50 (79%) female patients. A clamping method was used in which clamps were placed at L3 (46.6%), L4 (52.4%), and L2 in one case. There was a learning curve with this procedure, and 5 early cases had to be converted to an open procedure. Fifty-six (89%) of the patients had previously undergone some form of thoracic sympathectomy, and all had tried conservative measures. After a mean follow-up of 7 months, all patients considered their plantar hyperhidrosis symptoms to be “cured” or “improved;” 97% reported “cure.” All of the patients with previous thoracic sympathectomy had some degree of compensatory sweating. After lumbar sympathectomy, 51 of the 56 patients (91%) reported that their compensatory sweating was unchanged. In the 7 patients who did not have a previous thoracic sympathectomy, 1 reported mild and 6 reported moderate compensatory sweating. The authors stated that no sexual problems were reported by the male patients, and they did not discuss possible sexual problems among the female patients.

It is worth noting, that in contrast to earlier concerns about this procedure being associated with risks of permanent sexual dysfunction in men and women, the recent case series did not find any instances of permanent sexual dysfunction. A 2004 review from a multi-specialty working group on hyperhidrosis stated that lumbar sympathectomy is not recommended for plantar hyperhidrosis because of associated sexual dysfunction; this article did not cite any data documenting sexual dysfunction. (1) To date, there are very few studies on endoscopic lumbar sympathectomy for focal plantar hyperhidrosis and no comparative studies.

Conclusions: There are insufficient data supporting the safety and efficacy of lumbar sympathectomy for treating primary plantar hyperhidrosis.

Summary

There is insufficient evidence on the efficacy and safety of iontophoresis or microwave treatment for treating hyperhidrosis, and on radiofrequency ablation for palmar hyperhidrosis. There is evidence from randomized trials that botulinum toxin improves the net health outcome for patients with axillary hyperhidrosis and evidence that botulinum toxin A products improve the net health outcome for palmar hyperhidrosis. Due to the limited number of studies and high rates of adverse effects, there is insufficient evidence that botulinum toxin B improves the net health outcome for patients with primary palmar hyperhidrosis. There is insufficient evidence on the efficacy of any botulinum toxin products for other types of primary hyperhidrosis, including plantar and secondary hyperhidrosis.

Regarding surgical treatments for hyperhidrosis, data from randomized controlled trials and observational studies show high rates of efficacy of endoscopic transthoracic sympathectomy for primary focal hyperhidrosis, with the exception of plantar hyperhidrosis. There are, however, high rates of compensatory hyperhidrosis which must be considered in the treatment decision. There are insufficient data to draw conclusions on the efficacy of endoscopic lumbar sympathectomy in patients with primary plantar hyperhidrosis.

Practice Guidelines and Position Statements

In 2011, an expert consensus statement on the surgical treatment of hyperhidrosis was published by a task force of the Society of Thoracic Surgeons. (36) The document states that endoscopic thoracic sympathectomy is the treatment of choice for patients with primary hyperhidrosis. They further recommend the following treatment strategies (with R referring to rib and the number to which rib):

- R3 interruption for palmar hyperhidrosis; an R4 interruption is also reasonable. The authors note a slightly higher rate of compensatory sweating with an R3 but R3 is also more effective at treating hyperhidrosis.
- R4 or R5 interruption for palmar-axillary, palmar-axillary-plantar or axillary hyperhidrosis alone; R5 interruption is also an option for axillary hyperhidrosis alone.
- R3 interruption for craniofacial hyperhidrosis without blushing; an R2 and R3 procedure is an option but may lead to a higher rate of compensatory sweating, and also increases the risk of Horner's syndrome.

In 2008, the American Academy of Neurology created guidelines for use of botulinum neurotoxin for the treatment of autonomic disorders and pain. (37) These guidelines include the following recommendations for botulinum toxin injection as a treatment of hyperhidrosis:

- Should be offered as a treatment option to patients with axillary hyperhidrosis (Level A).
- Should be considered as a treatment option for palmar hyperhidrosis and drooling (Level B).
- May be considered for gustatory sweating (Level C).

CODING

The following codes for treatment and procedures applicable to this policy are included below for informational purposes. Inclusion or exclusion of a procedure, diagnosis or device code(s) does not constitute or imply member coverage or provider reimbursement. Please refer to the member's contract benefits in effect at the time of service to determine coverage or non-coverage of these services as it applies to an individual member.

CPT/HCPCS

32664 Thoracoscopy, surgical; with thoracic sympathectomy
 64650 Chemodenervation of eccrine glands; both axillae
 64653 Chemodenervation of eccrine glands; other area(s) (e.g., scalp, face, neck), per day
 69676 Tympanic neurectomy
 J0586 Injection, abobotulinumtoxinA, 5 units
 J0588 Injection, incobotulinumtoxinA, 1 unit

- A variety of iontophoretic devices can be purchased for use in the home. There are no specific HCPCS codes for these pieces of DME [durable medical equipment]. Code E1399 (Durable medical equipment, miscellaneous)
- Botulinum toxin may be coded for using HCPCS codes: J0585, J0586, J0587, J0588

ICD-9 Diagnoses

705.21 Primary focal hyperhidrosis
 705.22 Secondary focal hyperhidrosis
 780.8 Generalized hyperhidrosis

ICD-10 Diagnoses (Effective October 1, 2014)

L74.510 Primary focal hyperhidrosis, axilla
 L74.511 Primary focal hyperhidrosis, face
 L74.512 Primary focal hyperhidrosis, palms
 L74.513 Primary focal hyperhidrosis, soles
 L74.52 Secondary focal hyperhidrosis
 R61 Generalized hyperhidrosis

REVISIONS

09-23-2009	Policy added to the bcbsks.com web site as a stand-alone policy. Hyperhidrosis was previously discussed in the Botulinum Toxin medical policy.
01-26-2010	<p>In Policy Section:</p> <ul style="list-style-type: none"> ▪ Clarified policy language by changing 4.A. From: "Botulinum toxin when other treatments have failed" to " Botulinum toxin when symptoms are at a severity level of 3 or 4 (see severity scale below) and other treatments have failed" ▪ Clarified policy language by changing B.1. from: " Thoracic sympathectomy when symptoms are severe and other treatments have failed" to "Thoracic sympathectomy when symptoms are at a severity level of 3 or 4 (see severity scale below) and other treatments have failed"

03-31-2014	<p>Description section updated</p> <p>In Policy section:</p> <ul style="list-style-type: none"> ▪ Revised to create distinction between Primary Focal Hyperhidrosis and Secondary Hyperhidrosis. ▪ Updated Primary Focal Hyperhidrosis information From: "Treatment of hyperhidrosis resulting in significant impairment of daily activities may be considered medically necessary with the following: A. Non-surgical: 1. Topical treatments such as aluminum chloride, glutaraldehyde, anticholinergics, 2. Systemic treatments, such as anti-cholinergics, tranquilizers, 3. Iontophoresis, 4. Botulinum toxin when symptoms are at a severity level of 3 or 4 (see severity scale below) and other treatments have failed B. Surgical: 1. Thoracic sympathectomy when symptoms are at a severity level of 3 or 4 (see severity scale below) and other treatments have failed" To: "Treatment of primary hyperhidrosis may be considered medically necessary with any of the following medical complications: 1. acrocyanosis of the hands; or 2. history of recurrent skin maceration with bacterial or fungal infections; or 3. history of recurrent secondary infections; or 4. history of persistent eczematous dermatitis in spite of medical treatments with topical dermatological or systemic anticholinergic agents. <table border="1" data-bbox="376 932 1471 1703"> <thead> <tr> <th data-bbox="376 932 604 991">Focal Regions</th> <th data-bbox="604 932 1091 991">Treatments Considered Medically Necessary</th> <th data-bbox="1091 932 1471 991">Treatments Considered Investigational</th> </tr> </thead> <tbody> <tr> <td data-bbox="376 991 604 1251">Axillary</td> <td data-bbox="604 991 1091 1251"> <ul style="list-style-type: none"> ▪ aluminum chloride 20% solution* ▪ botulinum toxin* for severe primary axillary hyperhidrosis that is inadequately managed with topical agents*, in patients 18 years and older ▪ endoscopic transthoracic sympathectomy (ETS) and surgical excision of axillary sweat glands, if conservative treatment (i.e., aluminum chloride or botulinum toxin, individually and in combination) has failed </td> <td data-bbox="1091 991 1471 1251"> <ul style="list-style-type: none"> ▪ axillary liposuction ▪ iontophoresis ▪ microwave treatment </td> </tr> <tr> <td data-bbox="376 1251 604 1491">Palmar</td> <td data-bbox="604 1251 1091 1491"> <ul style="list-style-type: none"> ▪ aluminum chloride 20% solution* ▪ botulinum toxin A products for severe primary palmar hyperhidrosis that is inadequately managed with topical agents, in patients 18 years and older ▪ endoscopic transthoracic sympathectomy (ETS), if conservative treatment (i.e., aluminum chloride or botulinum toxin type A, individually and in combination) has failed </td> <td data-bbox="1091 1251 1471 1491"> <ul style="list-style-type: none"> ▪ RimabotulinumtoxinB ▪ iontophoresis ▪ microwave treatment ▪ radiofrequency ablation </td> </tr> <tr> <td data-bbox="376 1491 604 1591">Plantar</td> <td data-bbox="604 1491 1091 1591"> <ul style="list-style-type: none"> ▪ aluminum chloride 20% solution* </td> <td data-bbox="1091 1491 1471 1591"> <ul style="list-style-type: none"> ▪ botulinum toxin ▪ iontophoresis ▪ lumbar sympathectomy ▪ microwave treatment </td> </tr> <tr> <td data-bbox="376 1591 604 1703">Craniofacial</td> <td data-bbox="604 1591 1091 1703"> <ul style="list-style-type: none"> ▪ aluminum chloride 20% solution* ▪ endoscopic transthoracic sympathectomy (ETS), if conservative treatment (i.e., aluminum chloride) has failed </td> <td data-bbox="1091 1591 1471 1703"> <ul style="list-style-type: none"> ▪ botulinum toxin ▪ iontophoresis ▪ microwave treatment </td> </tr> </tbody> </table> <ul style="list-style-type: none"> ▪ Added section on indications for treatment of Secondary Gustatory Hyperhidrosis: "1.The following treatments would be considered medically necessary for the treatment of severe gustatory hyperhidrosis: a. aluminum chloride 20% solution* 	Focal Regions	Treatments Considered Medically Necessary	Treatments Considered Investigational	Axillary	<ul style="list-style-type: none"> ▪ aluminum chloride 20% solution* ▪ botulinum toxin* for severe primary axillary hyperhidrosis that is inadequately managed with topical agents*, in patients 18 years and older ▪ endoscopic transthoracic sympathectomy (ETS) and surgical excision of axillary sweat glands, if conservative treatment (i.e., aluminum chloride or botulinum toxin, individually and in combination) has failed 	<ul style="list-style-type: none"> ▪ axillary liposuction ▪ iontophoresis ▪ microwave treatment 	Palmar	<ul style="list-style-type: none"> ▪ aluminum chloride 20% solution* ▪ botulinum toxin A products for severe primary palmar hyperhidrosis that is inadequately managed with topical agents, in patients 18 years and older ▪ endoscopic transthoracic sympathectomy (ETS), if conservative treatment (i.e., aluminum chloride or botulinum toxin type A, individually and in combination) has failed 	<ul style="list-style-type: none"> ▪ RimabotulinumtoxinB ▪ iontophoresis ▪ microwave treatment ▪ radiofrequency ablation 	Plantar	<ul style="list-style-type: none"> ▪ aluminum chloride 20% solution* 	<ul style="list-style-type: none"> ▪ botulinum toxin ▪ iontophoresis ▪ lumbar sympathectomy ▪ microwave treatment 	Craniofacial	<ul style="list-style-type: none"> ▪ aluminum chloride 20% solution* ▪ endoscopic transthoracic sympathectomy (ETS), if conservative treatment (i.e., aluminum chloride) has failed 	<ul style="list-style-type: none"> ▪ botulinum toxin ▪ iontophoresis ▪ microwave treatment
Focal Regions	Treatments Considered Medically Necessary	Treatments Considered Investigational														
Axillary	<ul style="list-style-type: none"> ▪ aluminum chloride 20% solution* ▪ botulinum toxin* for severe primary axillary hyperhidrosis that is inadequately managed with topical agents*, in patients 18 years and older ▪ endoscopic transthoracic sympathectomy (ETS) and surgical excision of axillary sweat glands, if conservative treatment (i.e., aluminum chloride or botulinum toxin, individually and in combination) has failed 	<ul style="list-style-type: none"> ▪ axillary liposuction ▪ iontophoresis ▪ microwave treatment 														
Palmar	<ul style="list-style-type: none"> ▪ aluminum chloride 20% solution* ▪ botulinum toxin A products for severe primary palmar hyperhidrosis that is inadequately managed with topical agents, in patients 18 years and older ▪ endoscopic transthoracic sympathectomy (ETS), if conservative treatment (i.e., aluminum chloride or botulinum toxin type A, individually and in combination) has failed 	<ul style="list-style-type: none"> ▪ RimabotulinumtoxinB ▪ iontophoresis ▪ microwave treatment ▪ radiofrequency ablation 														
Plantar	<ul style="list-style-type: none"> ▪ aluminum chloride 20% solution* 	<ul style="list-style-type: none"> ▪ botulinum toxin ▪ iontophoresis ▪ lumbar sympathectomy ▪ microwave treatment 														
Craniofacial	<ul style="list-style-type: none"> ▪ aluminum chloride 20% solution* ▪ endoscopic transthoracic sympathectomy (ETS), if conservative treatment (i.e., aluminum chloride) has failed 	<ul style="list-style-type: none"> ▪ botulinum toxin ▪ iontophoresis ▪ microwave treatment 														

	<p>b. surgical options (i.e., tympanic neurectomy), if conservative treatment has failed.</p> <p>2. The following treatments are considered experimental / investigational as a treatment for severe gustatory hyperhidrosis including, but not limited to:</p> <p>a. botulinum toxin is considered investigational for treatment of gustatory hyperhidrosis. (See Policy Guidelines for list of conditions)</p> <p>b. iontophoresis</p> <p>*FDA approved indication (in the case of botulinum toxin, FDA approved for at least one botulinum toxin product)</p> <p>3. Treatment of hyperhidrosis is considered not medically necessary in the absence of functional impairment or medical complications."</p> <ul style="list-style-type: none"> ▪ Removed "Lumbar sympathectomy, axillary liposuction or resection of axillary sweat glands are considered experimental / investigational." ▪ Added Policy Guidelines leaving reference to the Severity Scale.
	Rationale section updated
	<p>In Coding section:</p> <ul style="list-style-type: none"> ▪ Added CPT and HCPCS codes: 69676, J0586, J0588 ▪ Revised nomenclature for HCPCS codes: J0585, J0587 ▪ Added ICD-10 Diagnosis Codes
	References updated

REFERENCES

1. Hornberger J, Grimes K, Naumann MM-SWGotR et al. Recognition, diagnosis, and treatment of primary focal hyperhidrosis. *J Am Acad Dermatol* 2004; 51(2):274-86.
2. Solish N, Bertucci V, Dansereau ACHAC et al. A comprehensive approach to the recognition, diagnosis, and severity-based treatment of focal hyperhidrosis: recommendations of the Canadian Hyperhidrosis Advisory Committee. *Dermatol Surg* 2007; 33(8):908-23.
3. Blue Cross and Blue Shield Association Technology Evaluation Center (TEC). Iontophoresis for Medical Indications. TEC Assessments 2003; Volume 18, Tab 3.
4. McAleer MA, Collins P. A Study Investigating Patients' Experience of Hospital and Home Iontophoresis for Hyperhidrosis. *J Dermatolog Treat* 2013 [Epub ahead of print].
5. Lowe NJ, Yamauchi PS, Lask GP et al. Efficacy and safety of botulinum toxin type A in the treatment of palmar hyperhidrosis: a double-blind, randomized, placebo-controlled study. *Dermatol Surg* 2002; 28(9):822-7.
6. Naumann M, Lowe NJ. Botulinum toxin type A in treatment of bilateral primary axillary hyperhidrosis: randomised, parallel group, double blind, placebo controlled trial. *BMJ* 2001; 323(7313):596-9.
7. Naumann MK, Hamm H, Lowe NJ. Effect of botulinum toxin type A on quality of life measures in patients with excessive axillary sweating: a randomized controlled trial. *Br J Dermatol* 2002; 147(6):1218-26.
8. Saadia D, Voustianiouk A, Wang AK et al. Botulinum toxin type A in primary palmar hyperhidrosis: randomized, single-blind, two-dose study. *Neurology* 2001; 57(11):2095-9.
9. Schnider P, Binder M, Auff E et al. Double-blind trial of botulinum A toxin for the treatment of focal hyperhidrosis of the palms. *Br J Dermatol* 1997; 136(4):548-52.
10. Talarico-Filho S, Mendonça DO, Nascimento M et al. A double-blind, randomized, comparative treatment of two type A botulinum toxins in the treatment of primary axillary hyperhidrosis. *Dermatol Surg* 2007; 33(1 Spec. No.):S44-S50.

11. Flanagan KH, King R, Glaser DA. Botulinum toxin type a versus topical 20% aluminum chloride for the treatment of moderate to severe primary focal axillary hyperhidrosis. *J Drugs Dermatol* 2008; 7(3):221-7.
12. Lowe NJ, Glaser DA, Eadie NNABiPAHCSG et al. Botulinum toxin type A in the treatment of primary axillary hyperhidrosis: a 52-week multicenter double-blind, randomized, placebo-controlled study of efficacy and safety. *J Am Acad Dermatol* 2007; 56(4):604-11.
13. Heckmann M, Ceballos-Baumann AO, Plewig G. Botulinum toxin A for axillary hyperhidrosis (excessive sweating). *N Engl J Med* 2001; 344(7):488-93.
14. Dressler D. Comparing Botox and Xeomin for axillar hyperhidrosis. *J Neural Transm* 2010; 117(3):317-9.
15. Baumann L, Slezinger A, Halem M et al. Pilot study of the safety and efficacy of Myobloc (botulinum toxin type B) for treatment of axillary hyperhidrosis. *Int J Dermatol* 2005; 44(5):418-24.
16. Baumann L, Slezinger A, Halem M et al. Double-blind, randomized, placebo-controlled pilot study of the safety and efficacy of Myobloc (botulinum toxin type B) for the treatment of palmar hyperhidrosis. *Dermatol Surg* 2005; 31(3):263-70.
17. Frasson E, Brigo F, Acler M et al. Botulinum toxin type A vs Type B for axillary hyperhidrosis in a case series of patients observed for 6 months. *Arch Dermatol* 2011; 147(1):122-3.
18. Glaser DA, Coleman WP, Fan LK et al. A randomized blinded clinical evaluation of a novel microwave device for treating axillary hyperhidrosis: the dermatologic reduction in underarm perspiration study. *Dermatol Surg* 2012; 38(2):185-91.
19. Hong HC, Lupin M, O'Shaughnessy KF. Clinical evaluation of a microwave device for treating axillary hyperhidrosis. *Dermatol Surg* 2012; 38(5):728-35.
20. Purtuloglu T, Atim A, Deniz S et al. Effect of radiofrequency ablation and comparison with surgical sympathectomy in palmar hyperhidrosis. *Eur J Cardiothorac Surg* 2013 [Epub ahead of print].
21. Clayman MA, Clayman SM, Seagle MB. A review of the surgical and medical treatment of Frey syndrome. *Ann Plast Surg* 2006; 57(5):581-4.
22. de Bree R, van der Waal I, Leeman CR. Management of Frey syndrome. *Head Neck* 2007; 29(8):773-8.
23. Hafner J, Beer GM. Axillary sweat gland excision. *Curr Probl Dermatol* 2002; 30:57-63.
24. Deng B, Tan QY, Jiang YG et al. Optimization of sympathectomy to treat palmar hyperhidrosis: the systematic review and meta-analysis of studies published during the past decade. *Surg Endosc* 2011; 25(6):1893-901.
25. Baumgartner FJ, Reyes M, Sarkisyan GG et al. Thoracoscopic sympathectomy for disabling palmar hyperhidrosis: A prospective randomized comparison between two levels. *Ann Thorac Surg* 2011; 92(6):2015-9.
26. Ishy A, de CJR, Wolosker N et al. Objective evaluation of patients with palmar hyperhidrosis submitted to two levels of sympathectomy: T3 and T4. *Interact Cardiovasc Thorac Surg* 2011; 12(4):545-8.
27. Gossot D, Galetta D, Pascal A et al. Long-term results of endoscopic thoracic sympathectomy for upper limb hyperhidrosis. *Ann Thorac Surg* 2003; 75(4):1075-9.
28. Han PP, Gottfried ON, Kenny KJ et al. Biportal thoracoscopic sympathectomy: surgical techniques and clinical results for the treatment of hyperhidrosis. *Neurosurgery* 2002; 50(2):306-12.
29. Leseche G, Castier Y, Thabut G et al. Endoscopic transthoracic sympathectomy for upper limb hyperhidrosis: limited sympathectomy does not reduce postoperative compensatory sweating. *J Vasc Surg* 2003; 37(1):124-8.

30. Lin TS, Kuo SJ, Chou MC. Uniportal endoscopic thoracic sympathectomy for treatment of palmar and axillary hyperhidrosis: analysis of 2000 cases. *Neurosurgery* 2002; 51(5 suppl):84-7.
31. Wait SD, Killory BD, Lekovic GP et al. Thoracoscopic sympathectomy for hyperhidrosis: analysis of 642 procedures with special attention to Horner's syndrome and compensatory hyperhidrosis. *Neurosurgery* 2010; 67(3):652-7.
32. Smidfelt K, Drott C. Late results of endoscopic thoracic sympathectomy for hyperhidrosis and facial blushing. *Br J Surg* 2011; 98(12):1719-24.
33. de Andrade Filho LO, Kuzniec S, Wolosker N et al. Technical difficulties and complications of sympathectomy in the treatment of hyperhidrosis: an analysis of 1731 cases. *Ann Vasc Surg* 2013 [Epub ahead of print].
34. Rieger R, Pedevilla S, Pochlauer S. Endoscopic lumbar sympathectomy for plantar hyperhidrosis. *Br J Surg* 2009; 96(12):1422-8.
35. Reifeld R. Endoscopic lumbar sympathectomy for focal plantar hyperhidrosis using the clamping method. *Surg Laparosc Endosc Percutan Tech* 2010; 20(4):231-6.
36. Cerfolio RJ, De CJR, Bryant AS et al. The Society of Thoracic Surgeons expert consensus for the surgical treatment of hyperhidrosis. *Ann Thorac Surg* 2011; 91(5):1642-8.
37. Naumann M, So Y, Argoff CE et al. Assessment: botulinum neurotoxin in the treatment of autonomic disorders and pain (an evidence-based review): report of the Therapeutics and Technology Assessment Subcommittee of the American Academy of Neurology. *Neurology* 2008; 70(19):1707-14.