

MEDICAL POLICY



SUBJECT: MINIMALLY INVASIVE/ MINIMAL ACCESS TECHNIQUES FOR LUMBAR INTERBODY FUSION	EFFECTIVE DATE: 08/20/09 REVISED DATE: 08/19/10, 09/15/11, 10/18/12, 09/19/13, 08/21/14
POLICY NUMBER: 7.01.83	PAGE: 1 OF: 10
CATEGORY: Technology Assessment	<ul style="list-style-type: none"><i>If the member's subscriber contract excludes coverage for a specific service it is not covered under that contract. In such cases, medical policy criteria are not applied.</i><i>Medical policies apply to commercial and Medicaid products only when a contract benefit for the specific service exists.</i><i>Medical policies only apply to Medicare products when a contract benefit exists and where there are no National or Local Medicare coverage decisions for the specific service.</i>

POLICY STATEMENT:

- I. Based upon our review and assessment of peer-reviewed literature, the following minimally invasive/minimal access techniques for interbody lumbar fusion have been medically proven to be effective and therefore can be considered as a **medically appropriate** treatment option to open standard lumbar fusion:
 - A. Anterior lumbar interbody fusion (ALIF);
 - B. Direct lateral interbody fusion (DLIF);
 - C. Extreme lateral interbody fusion (XLIF®);
 - D. Posterior lumbar interbody fusion (PLIF); or
 - E. Transforaminal lumbar interbody fusion (TLIF).
- II. Based upon our criteria and assessment of peer-reviewed literature, the following minimally invasive/minimal access techniques for interbody lumbar fusion have not been medically proven to be effective and are considered **investigational** either as a stand-alone procedure or as an adjunct to standard spinal fusion:
 - A. Axial lumbar interbody fusion (AxiaLIF®); or
 - B. Laparoscopic anterior lumbar interbody fusion (LALIF).

Refer to Corporate Medical Policy #11.01.03 regarding Experimental and Investigational Services.

POLICY GUIDELINES:

The Federal Employee Health Benefit Program (FEHBP/FEA) requires that procedures, devices or laboratory tests approved by the U.S. Food and Drug Administration (FDA) may not be considered investigational and thus these procedures, devices or laboratory tests may be assessed only on the basis of their medical necessity.

DESCRIPTION:

Lumbar fusion has become a widely accepted method for the management of a variety of disorders that require spinal stabilization, such as traumatic, degenerative, infectious, and neoplastic conditions. Interbody fusion of the lumbar spine can be approached from an anterior, posterior, and lateral direction. These approaches are traditionally performed with an open approach (long incision with wide retraction of the musculature. One of the drawbacks of conventional lumbar fusion is the extensive soft tissue dissection that is necessary in order to expose the anatomic landmarks for screw insertion, achieve a proper lateral-to-medial screw trajectory, and develop an acceptable fusion bed. The tissue injury that occurs during the surgical approach can result in increased postoperative pain, lengthened recovery time, and impaired spinal function. Blood loss during open lumbar fusion surgery can also be quite significant. These conventional approaches can now be performed through minimally invasive/minimal access procedures. A variety of minimally invasive/minimal access procedures are being investigated with the intent of limiting iatrogenic damage to muscular, ligamentous, neural, and vascular structures. Minimally invasive techniques that have been investigated include laparoscopic anterior lumbar fusion (LALIF), posterior lumbar interbody fusion (PLIF), transforaminal lumbar interbody fusion (TLIF), lateral interbody fusion (e.g., Extreme Lateral Interbody Fusion [XLIF] or Direct Lateral Interbody Fusion [DLIF]), and para-axial interbody fusion (AxiaLIF).

Anterior access provides direct visualization of the disc space through an abdominal incision, potentially allowing a more complete discectomy and better fusion than lateral or posterior approaches. An anterior approach avoids trauma to the paraspinal musculature, epidural scarring, traction on nerve roots, and dural tears. However, the retraction of the great

SUBJECT: MINIMALLY INVASIVE/ MINIMAL ACCESS TECHNIQUES FOR LUMBAR INTERBODY FUSION POLICY NUMBER: 7.01.83 CATEGORY: Technology Assessment	EFFECTIVE DATE: 08/20/09 REVISED DATE: 08/19/10, 09/15/11, 10/18/12, 09/19/13, 08/21/14 PAGE: 2 OF: 10
--	---

vessels, peritoneal contents, and superior hypogastric sympathetic plexus with a peritoneal or retroperitoneal approach place these structures at risk of iatrogenic injury. Access to the posterior space for the treatment of nerve compression is also limited. Laparoscopic Anterior Lumbar Interbody Fusion (LALIF) is a minimally invasive technique that has been proposed as an alternative to the open surgical approach to spinal fusion. This method employs a laparoscope to remove the diseased disc and insert an implant into the disc space intended to stabilize and promote fusion. This technique is evolving as a method of minimizing soft-tissue injury and is associated with a learning curve.

Posterior LIF can be performed through either a traditional open procedure with a midline incision or with a minimally invasive approach using bilateral paramedian incisions. In the open procedure, the midline muscle attachments are divided along the central incision to facilitate wide muscle retraction and laminectomy. Minimally invasive/minimal access PLIF uses tubular retractors (e.g., METRx™, Luxor™) to allow access and open visualization of the surgical area. These tubular retractors may be used to open smaller central bilateral working channels to access the pedicles and foramen. Minimally invasive PLIF typically involves partial laminotomies and facetectomies. The decompression allows treatment of spinal canal pathology (e.g., spinal stenosis, lateral recess and foraminal stenosis, synovial cysts, hypertrophic ligamentum flavum) as well as stabilization of the spine through interbody fusion.

Transforaminal LIF, performed through an open technique, is also performed through a posterior approach. Access to the spine is through the foramen which is enlarged by removal of surrounding bone. In minimally invasive TLIF, a single incision about 2-3 cm in length is made approximately 3 cm lateral to the midline. A tubular retractor is docked on the facet joint complex and a facetectomy with partial laminectomy is performed. Less dural retraction is needed with access through the foramen via unilateral facetectomy, and contralateral scar formation is eliminated. TLIF provides access to the posterior elements along with the intervertebral disc space.

Axial lumbar interbody fusion (AxiaLIF®, also called anterior para-axial, trans-sacral or paracoccygeal interbody fusion) is a minimally invasive technique designed to provide anterior access to the L4-S1 disc spaces for interbody fusion. It is performed percutaneously, under fluoroscopic guidance via the pre-sacral space. Theoretically, this approach avoids the viscera, blood vessels and nerves; preserves normal tissue at the treatment site; provides access to the disc space without interrupting the annulus; and allows for percutaneous longitudinal access to the anterior spine.

Lateral interbody fusion (e.g., Extreme Lateral Interbody Fusion [XLIF] or Direct Lateral Interbody Fusion [DLIF]) uses specialized retractors in a minimally invasive, lateral approach to the anterior spine through the psoas. In comparison with ALIF, the lateral approach does not risk injury to the peritoneum or great vessels. However, exposure to the spine may be more limited, and dissection of the psoas major places the nerves of the lumbar plexus at risk. Electromyographic monitoring and dissection predominantly within the anterior psoas major may be utilized to reduce the risk of nerve root injury. These various factors decrease the ability to perform a complete discectomy and address pathology of the posterior elements. The XLIF® surgical technique incorporates two systems developed by NuVasive®: the MaXcess® System and the NeuroVision® JJB System.

Both open and minimally invasive/minimal access interbody fusion surgeries may also include decompression of the spinal canal, use of interbody cages, bone grafts and osteoinductive agents (e.g., recombinant human bone morphogenetic protein), and insertion of pedicle screws and rods to increase stability of the spine.

RATIONALE:

Minimal access open anterior, posterior, and transforaminal LIF:

The available evidence (reviews, non-randomized comparative studies) suggests that after an initial training period, the mid-term health outcomes (including complication and fusion rates, pain and function) following minimally invasive anterior, posterior, transforaminal, and extreme lateral (XLIF) approaches are comparable to standard open approaches for single-level interbody fusion of the lumbar spine. Intra and peri-operative health outcomes (blood loss and hospital stay) have been shown to be improved (e.g., Kim, et al. 2010; Park, et al. 2007; Ghahreman, et al. 2010; Kasis, et al. 2009; Wang, et al. 2010; Wu, et al. 2010; Shunwu, et al. 2010; Rouben, et al. 2010) .

Direct lateral interbody fusion (DLIF):

DLIF procedure utilizes specialized FDA approved instrumentation from Medtronic. While well-designed, comparative clinical trials are needed to demonstrate whether these procedures provide improved health outcomes with long-term

SUBJECT: MINIMALLY INVASIVE/ MINIMAL ACCESS TECHNIQUES FOR LUMBAR INTERBODY FUSION POLICY NUMBER: 7.01.83 CATEGORY: Technology Assessment	EFFECTIVE DATE: 08/20/09 REVISED DATE: 08/19/10, 09/15/11, 10/18/12, 09/19/13, 08/21/14 PAGE: 3 OF: 10
--	---

follow-up, the outcomes from studies thus far demonstrate that DLIF has comparable outcomes to XLIF. P Berjano et al. (2012) conducted a retrospective cohort review of 97 consecutive patients from three centers with minimum 6-month follow-up (mean 12 months, 93 patients available for follow-up). The main diagnosis was DDD with or without stenosis, or spondylolisthesis, grade I. Functional status was evaluated by preoperative and last follow-up Oswestry Disability Index score. Leg and back pain were evaluated by visual analog scales. Complications were recorded and permanent complications and neurological impairment was actively investigated at last follow-up. Clinical success was considered to be achieved when the patient increased his functional ODI score by more than 12% or decreased his back pain VAS by more than 3 points. No permanent neurological impairment, vascular or visceral injuries were observed by the investigators. Transient neurological symptoms presented in 7% of cases, all resolved within 1 month from surgery. Transient thigh discomfort was observed in 9%. Clinical success was recorded in 92% of cases.

Extreme lateral interbody fusion (XLIF):

While extreme lateral interbody fusion as an endoscopic surgical procedure does not require FDA approval, the instrumentation associated with the XLIF procedure does. NuVasive® has developed the XLIF® instrumentation/ products for this surgical approach. This minimally invasive surgical platform is known as Maximum Access Surgery (MAS). MAS combines three categories of product offerings- NeuroVision®, MaXcess® and specialized implants such as SpheRx™ and CoRoent™. All surgical instrumentation associated with this procedure has received FDA approval either through the PMA or 510(k) process.

Ozgur et al. (2006) reported on the surgical technique for XLIF of the lower lumbar spine. 13 patients with axial low back pain who failed at least six months of conservative management underwent the XLIF technique. The authors concluded that, in comparison to anterior laparoscopic approaches, the XLIF approach had the advantages of not needing to retract the great vessels, not requiring a steep learning curve, and of no impairment to depth perception during the procedure. The most important advantage was a reduction in operative time. In this preliminary report, no complications were associated with the surgery; however, long-term follow-up and efficacy was yet to be reported.

In a 2009 report, Knight and colleagues compared complications from a series of 58 patients who underwent XLIF or DLIF (1- to 3-level) with a historical cohort of patients who underwent open posterolateral lumbar fusion. Thirteen patients (22.4%) experienced a mild or major complication. Nine of the complications were approach-related (2 L4 nerve root injuries, 6 cases of meralgia paresthetica, and 1 case of significant psoas muscle spasm). In 4 additional cases, the procedure was aborted because of concerns about nerve proximity. Compared with the historical cohort, there was less blood loss (136 vs. 489 mL), a shorter operative time (161 vs. 200 mins.), similar hospital stay (5 days), and a similar percentage of complications (22.4 vs. 22.5%). Approach-related complications in the open cohort included wound infection and dural tears.

In 2010, Rodgers et al. published a retrospective review of a database for all patients treated with the XLIF procedure by a single surgeon (between 2006 and 2008), focusing on early complications (less than 3 months) in obese and nonobese patients. Out of a total of 432 patients treated with XLIF during this period, 313 (72%) met the inclusion criteria for the study and had complete data; 156 were obese (greater than 30 kg/m²) and 157 were not obese. Patients who were obese were slightly younger (58.9 vs. 62.9 years of age) and had a higher incidence of diabetes mellitus (48 vs. 17) than patients who were not obese, but were otherwise comparable at baseline. There were 27 complications (8.6%) in the entire group, which included cardiac and wound complications, vertebral body fractures (1 requiring reoperation), nerve injuries, gastrointestinal injuries (1 requiring reoperation), and hardware failures (1 requiring reoperation for recurrent stenosis after cage subsidence). The complication and reoperation rates were not significantly different between the obese and nonobese groups. There were no cerebrospinal fluid leaks, no infections, and no patient required transfusion. The average length of hospital stay was 1.2 days. The authors noted that reliable automated neurological monitoring and fluoroscopic guidance, and meticulous attention to operative technique are required, but that the early outcomes compare well with traditional interventions.

In 2011, Rodgers and colleagues reported a retrospective analysis of intraoperative and perioperative complications from all consecutive patients (600 procedures, 741 levels) treated by 2 surgeons since the XLIF procedure was introduced at their institution. Four-hundred eighty-five procedures were single level, 90 were 2 level, and 25 involved 3 or more levels. The hospital stay averaged 1.2 days. There were 37 complications (6%), classified into medical (60%) and surgical

SUBJECT: MINIMALLY INVASIVE/ MINIMAL ACCESS TECHNIQUES FOR LUMBAR INTERBODY FUSION POLICY NUMBER: 7.01.83 CATEGORY: Technology Assessment	EFFECTIVE DATE: 08/20/09 REVISED DATE: 08/19/10, 09/15/11, 10/18/12, 09/19/13, 08/21/14 PAGE: 4 OF: 10
--	---

(40%). Surgical complications included 4 transient postoperative neurologic deficits and 1 subcutaneous hematoma. There were no wound infections, no vascular injuries, and no intraoperative visceral injuries in this series. At a minimum 1-year follow-up, VAS pain scores had decreased from an average 8.8 to 3.1.

Laparoscopic anterior interbody lumbar fusion:

Currently, the published, peer-reviewed scientific literature does not allow strong conclusions regarding the overall benefit and long-term efficacy of the laparoscopic anterior approach compared to open spinal fusion. Studies also report a potentially a higher rate of complications with laparoscopic ALIF.

In review of the literature on laparoscopic ALIF, Inamasu et al, (2005) identified 19 studies which described the outcome of a L5-S1 laparoscopic ALIF, 9 studies which described the outcome of the L4-L5 laparoscopic ALIF, and 8 studies which described the outcome of a 2-level laparoscopic ALIF. The review concluded that there was no marked difference between laparoscopic ALIF and the open or mini-open ALIF in terms of short-term efficacy (operative time, blood loss, and length of hospital stay), but there was a higher incidence of complications. In addition, the conversion rate to open surgery was considered to be high. It was noted that at the time of the review article, some spine surgeons were abandoning the laparoscopic approach and switching to mini-open ALIF.

The largest trial on laparoscopic ALIF was a prospective multicenter (19 surgeons from 10 U.S. centers) investigational device exemption (FDA-regulated) trial, published in 1999 by Regan, et al, that compared short-term outcomes from laparoscopic fusion of the spine (240 consecutive patients) and open ALIF (earlier cohort of 591 similar patients). Inclusion criteria were painful degenerative disc disease consisting of disc space narrowing at 1 or 2 contiguous levels (L4-L5 and L5-S1). Single level fusion was performed in 215 patients using laparoscopy and in 305 patients using the open procedure; 2-level fusions were performed in 25 patients via laparoscopy and 286 patients with the open procedure. In 25 (10%) of the laparoscopy patients, conversion to an open procedure was required due to bleeding (n=6), anatomic considerations (n=5), adhesions or scar tissue limiting access to the spine (n=8); and technical difficulties in placing the threaded cage (n=6). The hospital stay was modestly shorter for the single-level laparoscopy group (3.3 vs. 4 days), but not for patients undergoing 2-level laparoscopy. Operative time was increased (201 vs. 142 minutes) for the single-level laparoscopic approach (243 minutes for the 25 cases converted to open). For 2-level laparoscopy, the procedure time was 146 minutes longer than for the open approach. The reoperation rate for single-level procedures was 4.7% in the laparoscopy group compared with 2.3% in the open group (not significantly different). Major complications (implant migration, great vessel damage, pulmonary embolism) were significantly lower in the laparoscopy group (0% vs. 2%). Postoperative complications were similar in the 2 groups, with an occurrence of 14.1% in the open approach and 19.1% for the laparoscopic approach.

A prospective comparison of 50 consecutive patients (25 in each group) with disabling discogenic pain who underwent 1 or 2 level ALIF at L4-L5 with either a laparoscopic or mini-open approach was reported by Zdeblick and David in 2000. There was no difference between the laparoscopic and mini-open approaches in operating time (125 vs. 123 minutes), blood loss (50 cc vs. 55 cc), or length of hospital stay (1.4 vs. 1.3 days) for single-level fusion. For 2-level fusion, the operating time was increased for the laparoscopic procedure (185 vs. 160 minutes). There was a 20% rate of complications in the laparoscopic group (disc herniation, ureter injury, iliac vein laceration, transient retrograde ejaculation, deep vein thrombosis) compared with 4% in the mini-open group (ileus). Exposure was considered inadequate in the laparoscopic group, with only a single interbody cage placed in 16% of patients in the laparoscopic group. All patients in the mini-open group had 2 interbody cages placed.

AxiaLIF:

The AxiaLIF and AxiaLIF 2 Level Systems were developed by TranS1 and consist of techniques and surgical instruments for creating a pre-sacral access route to perform percutaneous fusion of the L5 - S1 or L4 - S1 vertebral bodies. The AxiaLIF 2 level system received premarket notification in April 2008. FDA premarket notification [510(k)] summaries indicate that the procedures are intended to provide anterior stabilization of the spinal segments as an adjunct to spinal fusion and for assisting in the treatment of degeneration of the lumbar disc, performing lumbar discectomy, or for assistance in the performance of interbody fusion.

There is insufficient evidence to determine if axial lumbar interbody fusion is as effective or as safe as other established surgical techniques.

SUBJECT: MINIMALLY INVASIVE/ MINIMAL ACCESS TECHNIQUES FOR LUMBAR INTERBODY FUSION POLICY NUMBER: 7.01.83 CATEGORY: Technology Assessment	EFFECTIVE DATE: 08/20/09 REVISED DATE: 08/19/10, 09/15/11, 10/18/12, 09/19/13, 08/21/14 PAGE: 5 OF: 10
--	---

Aryan and colleagues report on their series of 35 patients with average follow-up of 17.5 months. These patients had pain secondary to lumbar degenerative disc disease, degenerative scoliosis, or lytic spondylolisthesis. In 21 of the patients, the AxiaLIF procedure was followed by percutaneous pedicle screw-rod fixation, 2 patients had extreme lateral interbody fusion combined with posterior instrumentation, and 10 had a stand-alone procedure. Two patients had axial LIF as part of a larger construct after unfavorable anatomy prevented access to the L5-S1 disc space during open lumbar fusion. Thirty-two patients had radiographic evidence of stable cage placement and fusion at last follow-up.

In 2010, Patil and colleagues reported a retrospective review of 50 patients treated with AxiaLIF. Four patients (8%) underwent 2-level AxiaLIF and 16 patients (32%) underwent a combination of AxiaLIF with another procedure for an additional level of fusion. There were 3 reoperations due to pseudoarthrosis (n=2) and rectal injury (n=1). Other complications included superficial infection (n=5), hematoma (n=2), and irritation of a nerve root by a screw (n=1). At 12- to 24-month follow-up VAS scores had decreased from 8.1 to 3.6 (n = 48). At an average 12-month follow-up, 47 of 49 patients (96%) with postoperative radiographs achieved solid fusion. There were no significant differences between pre- and postoperative disk space height and lumbar lordosis angle.

CODES: Number Description

Eligibility for reimbursement is based upon the benefits set forth in the member's subscriber contract.

CODES MAY NOT BE COVERED UNDER ALL CIRCUMSTANCES. PLEASE READ THE POLICY AND GUIDELINES STATEMENTS CAREFULLY.

Codes may not be all inclusive as the AMA and CMS code updates may occur more frequently than policy updates.

Code Key: Experimental/Investigational = (E/I), Not medically necessary/ appropriate = (NMN).

CPT: Minimally invasive/minimal access ALIF, PLIF or TLIF would be billed using open lumbar fusion/arthrodesis codes.

22586 (E/I)	Arthrodesis, pre-sacral interbody technique, including disc preparation, discectomy, with posterior instrumentation with image guidance, includes bone graft with preformed, L5-S1 interspace
0195T (E/I)	Arthrodesis, pre-sacral interbody technique, including disc preparation, discectomy, without instrumentation with image guidance, includes bone graft with preformed, L5-S1 interspace
0196T (E/I)	L4-L5 interspace (list separately in addition to code for primary)
0309T (E/I)	Arthrodesis, pre-sacral interbody technique, including disc preparation, discectomy, with posterior instrumentation with image guidance, includes bone graft with preformed, L4-L5 interspace (list separately in addition to code for primary procedure)

No specific codes exist for billing of DLIF or XLIF

Copyright © 2014 American Medical Association, Chicago, IL

HCPCS: No specific codes

ICD9: Multiple diagnosis codes

ICD10: Multiple diagnosis codes

REFERENCES:

Acosta FL, et al. Changes in coronal and sagittal plane alignment following minimally invasive direct lateral interbody fusion for the treatment of degenerative lumbar disease in adults: a radiographic study. *J Neurosurg Spine* 2011 Apr 8 [Epub ahead of print].

Aichmair A, et al. An institutional six-year trend analysis of the neurological outcome after lateral interbody fusion: 1 6-year trend analysis of a single institution. *Spine* 2013 Nov 1;38(23):E1483-90.

SUBJECT: MINIMALLY INVASIVE/ MINIMAL ACCESS TECHNIQUES FOR LUMBAR INTERBODY FUSION POLICY NUMBER: 7.01.83 CATEGORY: Technology Assessment	EFFECTIVE DATE: 08/20/09 REVISED DATE: 08/19/10, 09/15/11, 10/18/12, 09/19/13, 08/21/14 PAGE: 6 OF: 10
--	---

*Anand N, et al. Minimally invasive multilevel percutaneous correction and fusion for adult lumbar degenerative scoliosis: a technique and feasibility study. *J Spinal Disord Tech* 2008 Oct;21(7):459-67.

Arnold PM, et al. The lateral transpsoas approach to the lumbar and thoracic spine: A review. *Surg Neurol Int* 2012;3(Suppl 3):S198-215.

*Aryan HE, et al. Percutaneous axial lumbar interbody fusion (AxiaLIF) of the L5-S1 segment: initial clinical and radiographic experience. *Min Invasive Neurosurg* 2008 Aug;51(4):225-30.

Berjano P, et al. Direct lateral access lumbar and thoracolumbar fusion: preliminary results. *Eur Spine J* 2012 May;21 Suppl 1:S37-42.

Berjano P, et al. Far lateral approaches (XLIF) in adult scoliosis. *Eur Spine J* 2013 Mar;22 Suppl 2:S242-53.

*BlueCross BlueShield Association. Minimally invasive lumbar interbody fusion - archived. Medical Policy Reference Manual #7.01.115. archived 2011 Oct 11.

BlueCross BlueShield Association. Axial lumbosacral interbody fusion. Medical Policy Reference Manual #7.01.130. 2013 Nov 14.

Boachie-Adjei O, et al. Axial lumbar interbody fusion (AxiaLIF) approach for adult scoliosis. *Eur Spine J* 2013 Mar Suppl 2:S225-31.

Caputo AM, et al. Extreme lateral interbody fusion for the treatment of adult degenerative scoliosis. *J Clin Neurosci* 2013 Nov;20(11):1558-63.

Dakwar E, et al. Early outcomes and safety of the minimally invasive, lateral retroperitoneal transpsoas approach for adult degenerative scoliosis. *Neurosurg Focus* 2010 Mar;28(3):E8.

*Eck JC, et al. Minimally invasive lumbar spinal fusion. *J Am Acad Orthop Surg* 2007 Jun;15(6):321-9.

Elowitz EH, et al. Evaluation of indirect decompression of the lumbar spinal canal following minimally invasive lateral transpsoas interbody fusion: radiographic and outcome analysis. *Minim Invasive Neurosurg* 2011 Oct;54(5-6):201-6.

*Foley KT, et al. Minimally invasive lumbar fusion. *Spine* 2003 Aug 1;28(15 Suppl):S26-35.

Fourney DR, et al. Does minimal access tubular assisted spine surgery increase or decrease complications in spinal decompression or fusion? *Spine* 2010 Apr 20;35(9 Suppl):S57-65.

Gerszten PC, et al. Retrospective analysis of L5-S1 axial lumbar interbody fusion (AxiaLIF): a comparison with and without the use of recombinant human bone morphogenetic protein-2. *Spine J* 2011 Nov;11(11):1027-32.

Gerszten PC, et al. Axial presacral lumbar interbody fusion and percutaneous posterior fixation for stabilization of lumbosacral isthmic spondylolisthesis. *J Spinal Disord Tech* 2012 Apr;25(2):E36-40.

Ghahreman A, et al. Minimal access versus open posterior lumbar interbody fusion in the treatment of spondylolisthesis. *Neurosurg* 2010 Feb;66(2):296-304.

Gundanna M, et al. Complications with axial presacral lumbar interbody fusion : A 5-year postmarketing surveillance experience. *SAS Journal* 2011 Sep;5:90-4.

Hadjipavlou A, et al. Safety and effectiveness of retrorectal presacral approach for lumbosacral axial instrumentation. A clinical study. *Acta Orthop* 2013 Apr;79(2):220-9.

*Inamasu J, et al. Laparoscopic anterior lumbar interbody fusion: a review of outcome studies. *Minim Invasive Neurosurg* 2005 Dec;48(6):340-7.

Isaacs RE, et al. A prospective, nonrandomized, multicenter evaluation of extreme lateral interbody fusion for the treatment of adult degenerative scoliosis: perioperative outcomes and complications. *Spine* 2010 Dec 15;35(26 Suppl):S322-30.

SUBJECT: MINIMALLY INVASIVE/ MINIMAL ACCESS TECHNIQUES FOR LUMBAR INTERBODY FUSION POLICY NUMBER: 7.01.83 CATEGORY: Technology Assessment	EFFECTIVE DATE: 08/20/09 REVISED DATE: 08/19/10, 09/15/11, 10/18/12, 09/19/13, 08/21/14 PAGE: 7 OF: 10
--	---

Karikari IO, et al. Minimally invasive lumbar interbody fusion in patients over seventy years of age: analysis of peri-and post-operative complications. Neurosurg 2011 Jan 6 [Epub ahead of print].

Karikari IO, et al. Minimally invasive transforaminal lumbar interbody fusion: a review of techniques and outcomes. Spine 2010 Dec 15;35(26 Suppl):S294-301.

Karikari IO, et al. Extreme lateral interbody fusion approach for isolated thoracic and thoracolumbar spine disease: initial clinical experience and early outcomes. J Spinal Disord Tech 2011 Aug;24(6):368-375.

Kasis AG, et al. Significantly improved outcomes with less invasive posterior lumbar interbody fusion incorporating total facetectomy. Spine 2009 Mar 15;34(6):572-7.

Kepler CK, et al. Lateral transpsoas interbody fusion (LTIF) with plate fixation and unilateral pedicle screws: a preliminary report. J Spinal Disord Tech 2010 Dec 8 [Epub ahead of print].

Kim JS, et al. Minimally invasive anterior lumbar interbody fusion followed by percutaneous pedicle screw fixation for isthmic spondylolisthesis: minimum 5-year follow-up. Spine J 2010 May;10(5):404-9.

Kim JS, et al. Which lumbar interbody fusion technique is better in terms of level for the treatment of unstable isthmic spondylolisthesis? J Neurosurg Spine 2010 Feb;12(2):171-7.

Kim JS, et al. Mini-transforaminal lumbar interbody fusion versus anterior lumbar interbody fusion augmented by percutaneous pedicle screw fixation: a comparison of surgical outcomes in adult low-grade isthmic spondylolisthesis. J Spinal Disord Tech 2009 Apr;22(2):114-21.

Knight RQ, et al. Direct lateral lumbar interbody fusion for degenerative conditions: early complication profile. J Spinal Disord Tech 2009 Feb;22(1):34-7.

*Lee SH, et al. Minimally invasive anterior lumbar interbody fusion followed by percutaneous pedicle screw fixation for isthmic spondylolisthesis. Spine J 2004 Nov-Dec;4(6):644-9.

Lee JC, et al. Learning curve and clinical outcomes of minimally invasive transforaminal lumbar interbody fusion: our experience in 86 consecutive cases. Spine 2012 Aug 15;37(18):1548-57.

Lee KH, et al. Clinical and radiological outcomes of open versus minimally invasive transforaminal lumbar interbody fusion. Eur Spine J 2012 Nov;21(11):2265-70.

Lindley EM, et al. Complications of axial lumbar interbody fusion. J Neurosurg Spine 2011 Sep;15(3):273-9.

Lykissas MG, et al. Is there any relation between the amount of curve correction and postoperative neurological deficit or pain in patients undergoing standalone lateral lumbar interbody fusion? Spine 2013 Sep 1;38(19):1656-62.

Lykissas MG, et al. Nerve injury after lateral lumbar interbody fusion: a review of 919 treated levels with identification of risk factors. Spine J 2014 May 1;14(5):749-58.

Lykissas MG, et al. Nerve injury and recovery after lateral lumbar interbody fusion with and without bone morphogenetic protein-2 augmentation: a cohort-controlled study. Spine J 2014 Feb 1;14(2):217-24.

*MacMillan M. Computer-guided percutaneous interbody fixation and fusion of the L5-S1 disc: a 2-year prospective study. J Spinal Disord Tech 2005 Feb;18 Suppl:S90-5.

Malham GM, et al. Clinical outcomes and fusion rates after the first 30 extreme lateral interbody fusions. Scientific World J 2012;2012:246989.

Malham GM, et al. maintenance of segmental lordosis and disc height in standalone and instrumented extreme lateral interbody fusion (XLIF). J Spinal Disord Tech 2014 Mar 24 [Epub ahead of print].

Marchi L, et al. Results and complications after 2-level axial lumbar interbody fusion with a minimum 2-year follow-up. J Neurosurg Spine 2012 Sep;1793:187-92.

SUBJECT: MINIMALLY INVASIVE/ MINIMAL ACCESS TECHNIQUES FOR LUMBAR INTERBODY FUSION POLICY NUMBER: 7.01.83 CATEGORY: Technology Assessment	EFFECTIVE DATE: 08/20/09 REVISED DATE: 08/19/10, 09/15/11, 10/18/12, 09/19/13, 08/21/14 PAGE: 8 OF: 10
--	---

National Institute for Health and Clinical Excellence (NICE). Lateral (including extreme, extra and direct lateral) interbody fusion in the lumbar spine. IPG 321. Nov 2009 [<http://www.nice.org.uk/guidance/index>] accessed 7/16/14.

National Institute for Health and Clinical Excellence (NICE). Transaxial interbody lumbosacral fusion. IPG 387. 2011 Mar [<http://www.nice.org.uk/guidance/index>] accessed 7/15/14.

Ntoukas V, et al. Minimally invasive approach versus traditional open approach for one level posterior lumbar interbody fusion. Minim Invasive Neurosurg 2010 Feb;53(1):21-4.

Oliveira L, et al. A radiographic assessment of the ability of the extreme lateral interbody fusion procedure to indirectly decompress the neural elements. Spine 2010 Dec 15;35(26 Suppl):S331-7.

Oppenheimer JH, et al. Minimally invasive spine technology and minimally invasive spine surgery: a historical review. Neurosurg Focus 2009 Sep;27(3):E9.

*Ozgur BM, et al. Combined XLIF and AXiaLIF techniques for minimally-invasive surgical access. Internet J Minim Invasive Spinal Tech 2008;2(3).

*Park Y, et al. Comparison of one-level posterior lumbar interbody fusion performed with a minimally invasive approach or a traditional open approach. Spine 2007 Mar 1;32(5):537-43.

Parker SL, et al. Cost effectiveness of minimally invasive versus open transforaminal lumbar interbody fusion for degenerative spondylolisthesis associated low-back pain and leg pain over two years. World Neurosurg 2012 Jul;78(1-2):178-84.

Parker SL, et al. Minimally invasive versus open transforaminal lumbar interbody fusion (TLIF) for degenerative spondylolisthesis: comparative effectiveness and cost-utility analysis. World Neurosurg 2013 Jan 12 [Epub ahead of print].

Patil SS, et al. Clinical and radiological outcomes of axial lumbar interbody fusion. Orthopedics 2010 Dec 1; 33(2):883.

Phillips FM, et al. Adult degenerative scoliosis treated with XLIF: clinical and radiographical results of a prospective multicenter study with 24-month follow-up. Spine 2013 Oct 1;38(21):1853-61.

*Regan JJ, et al. Laparoscopic approach to L4-L5 for interbody fusion using BAK cages: experience in the first 58 cases. Spine 1999 Oct 15;24(20):2171-4.

Regev GJ, et al. Incidence and prevention of intervertebral cage overhang with minimally invasive lateral approach fusions. Spine 2010 Jun 15;35(14):1406-11.

*Rodgers WB, et al. Extreme lateral interbody fusion in the morbidly obese. Spine J 2008 Sep-Oct;8(5 Suppl 1):10S.

Rodgers WB, et al. Early complications of extreme lateral interbody fusion in the obese. J Spinal Disord Tech 2010 Jan 15.[Epub ahead of print].

Rodgers WB, et al. Grade 2 spondylolisthesis at L4-5 treated by XLIF: safety and midterm results in the “worst case scenario”. Scientific World J 2012;2012:356712.

Rodgers WB, et al. Intraoperative and early postoperative complications in extreme lateral interbody fusion: an analysis of 600 cases. Spine 2011 Jan 1;36(1):26-32.

Rouben D, et al. Long-term durability of minimal invasive posterior transforaminal lumbar interbody fusion: a clinical and radiographic follow-up. J Spinal Disord Tech 2010 Oct 21 [Epub ahead of print].

*Shen FH, et al. Minimally invasive techniques for lumbar interbody fusions. Orthop Clin North Am 2007 Jul;38(3):373-86.

Shunwu F, et al. Minimally invasive transforaminal lumbar interbody fusion for the treatment of degenerative lumbar diseases. Spine 2010 Aug 1;35(17):1615-20.

SUBJECT: MINIMALLY INVASIVE/ MINIMAL ACCESS TECHNIQUES FOR LUMBAR INTERBODY FUSION POLICY NUMBER: 7.01.83 CATEGORY: Technology Assessment	EFFECTIVE DATE: 08/20/09 REVISED DATE: 08/19/10, 09/15/11, 10/18/12, 09/19/13, 08/21/14 PAGE: 9 OF: 10
--	---

Smith W, et al. Complications in extreme lateral interbody fusion: One surgeon's experience with 566 patients and 1550 levels. Presented at: 24th annual meeting of the North American Spine Society (NASS). Abstract P124. Spine J 2009;9:178S-9S.

Smith WD, et al. A comparison of perioperative changes and outcome between open and mini-open approaches for anterior lumbar discectomy and fusion. J Clin Neurosci 2012 May;19(5):673-80.

Society for Minimally Invasive Spine Surgery. SMISS Position Statement on Presacral Lumbar Interbody Fusion. 2012 Feb [http://www.smiss.org/] accessed 7/16/14.

*Stippler M, et al. Outcomes after percutaneous TranS1 AxiaLIF® L5-S1 interbody fusion for intractable lower back pain. Internet J Neurosurg 2008;5(1).

Tender GC, et al. Percutaneous pedicle screw reduction and axial presacral lumbar interbody fusion for treatment of lumbosacral spondylolisthesis: A case series. J Med Case Rep 2011 Sep 12;5:454.

Tobler WD, et al. Minimally- invasive axial presacral L5-S1 interbody fusion: Two-year clinical and radiographic outcomes. Spine 2011 Sep 15;36(20):E1296-301.

Tobler WD, et al. The presacral retroperitoneal approach for axial lumbar interbody fusion: a prospective study of clinical concerns, complications and fusion rates at a follow-up of two years in 26 patients. J Bone Joint Surg Br 2011 Jul;93(7):955-60.

Tobler WD, et al. Clinical and radiographic outcomes with L4-S1 axial lumbar interbody fusion (AxiaLIF) and posterior instrumentation: a multicenter study. Med Devices 2013 Sep 18;6:155-61.

Tormenti MJ, et al. Complications and radiographic correction in adult scoliosis following combined transpsoas extreme lateral interbody fusion and posterior pedicle screw instrumentation. Neurosurg Focus 2010 Mar;28(3):E7.

Wang J, et al. Comparison of one-level minimally invasive and open transforaminal lumbar interbody fusion in degenerative and isthmic spondylolisthesis grades 1 and 2. Eur Spine J 2010 Oct;19(10):1780-4.

Wang J, et al. Minimally invasive or open transforaminal lumbar interbody fusion as revision surgery for patients previously treated by open discectomy and decompression of the lumbar spine. Eur Spine J 2011 Apr;20(4):623-8.

Wang J, et al. Comparison of clinical outcome in overweight or obese patients after minimally invasive versus open transforaminal lumbar interbody fusion. J Spinal Disord Tech 2012 May 17 [Epub ahead of print].

Whang PG, et al. Comparison of axial and anterior interbody fusions of the L5-S1 segment: a retrospective cohort analysis. J Spinal Disord Tech 2013 Dec;26(8):437-43.

Wu RH, et al. Minimal access versus open transforaminal lumbar interbody fusion: a meta-analysis of fusion rates. Spine 2010 Dec 15;35(26):2273-81.

Youssef JA, et al. Minimally invasive surgery: lateral approach interbody fusion: results and review. Spine 2010 Dec 15;35(26 Suppl):S302-1.

*Zdeblick TA, et al. A prospective comparison of surgical approach for anterior L4-L5 fusion: laparoscopic versus mini anterior lumbar interbody fusion. Spine 2000 Oct 15;25(20):2682-7.

Zeilstra DJ, et al. Axial lumbar interbody fusion: 6-year single-center experience. Clin Interv Aging 2013;8:1063-9.

* key article

KEY WORDS:

Axial, Direct Lateral Interbody Fusion, Extracavitary, Extreme lateral, Interbody fusion, Laparoscopic anterior, Minimal access, Paracoccygeal axial approach, Pre-sacral approach, Trans-sacral approach

SUBJECT: MINIMALLY INVASIVE/ MINIMAL ACCESS TECHNIQUES FOR LUMBAR INTERBODY FUSION	EFFECTIVE DATE: 08/20/09
POLICY NUMBER: 7.01.83	REVISED DATE: 08/19/10, 09/15/11, 10/18/12, 09/19/13, 08/21/14
CATEGORY: Technology Assessment	PAGE: 10 OF: 10

CMS COVERAGE FOR MEDICARE PRODUCT MEMBERS

Based upon our review, minimally invasive/minimal access lumbar interbody fusion is not specifically addressed in National or Regional Medicare coverage determinations/policies. However, there is currently a Local Coverage Determination (LCD) and related article for Category III CPT codes. Please refer to the following LCD website for Medicare Members:

http://apps.ngsmedicare.com/lcd/LCD_L25275.htm
http://apps.ngsmedicare.com/SIA/ARTICLE_A46075.htm