

SUBJECT: PLUGS FOR FISTULA REPAIR

EFFECTIVE DATE: 08/18/11

POLICY NUMBER: 7.01.86

REVISED DATE: 07/19/12, 06/20/13, 05/22/14

CATEGORY: Technology Assessment

PAGE: 1 OF: 6

- *If the member's subscriber contract excludes coverage for a specific service it is not covered under that contract. In such cases, medical policy criteria are not applied.*
- *Medical policies apply to commercial and Medicaid products only when a contract benefit for the specific service exists.*
- *Medical policies only apply to Medicare products when a contract benefit exists and where there are no National or Local Medicare coverage decisions for the specific service.*

POLICY STATEMENT:

Based upon our criteria and assessment of peer-reviewed literature, biosynthetic fistula plugs, including plugs made of porcine small intestine submucosa or of synthetic material, have not been medically proven effective and are considered **investigational** for all indications including, but not limited to, the repair of anal and rectal fistulas.

POLICY GUIDELINES:

The Federal Employee Health Benefit Program (FEHBP/FEP) requires that procedures, devices or laboratory tests approved by the U.S. Food and Drug Administration (FDA) may not be considered investigational and thus these procedures, devices or laboratory tests may be assessed only on the basis of their medical necessity.

DESCRIPTION:

An anal fistula is an abnormal communication between the interior of the anal canal or rectum and the skin surface. Rarer forms may communicate with the vagina or other pelvic structures, including the bowel. Most fistulas begin as anorectal abscesses. When the abscess opens spontaneously into the anal canal (or has been opened surgically), a fistula may occur. Anal fistulas are described as low (present in the lower part and not extending up to the anorectal sling) or high (extending up to or beyond the anorectal sling). High fistula can be associated with incontinence. Anal fistulas are also classified according to their relationship with the external sphincter. Intersphincteric fistulas are the most common and cross only the internal sphincter. Transsphincteric fistulas pass through the internal and external sphincters. The type of surgical treatment depends on the location and complexity of the fistula. Treatments include fistulotomy/fistulectomy, endorectal/anal sliding flaps, seton drain, and fibrin glue. Lay-open fistulotomy in high fistulas carries risk of incontinence. Draining setons can control sepsis but few patients heal after removal of the seton and they are poorly tolerated long term. Cutting setons can cause continence disturbances. Because of recurrence rates and the significant risk of incontinence with these surgical procedures, sphincter-preserving techniques such as fistula plugs have been evaluated and proposed as an alternative method in the treatment of anorectal fistulas.

Anal fistula plugs are biosynthetic devices used to promote healing and prevent recurrence of an anal fistula. In a minimally invasive procedure, the fistula tract is identified using a probe or imaging techniques and then cleaned by irrigation. The conical-shaped fistula plug is pulled into the tract until it blocks the internal opening and then is anchored in place with sutures. The external opening is not completely sealed so that drainage of the fistula can continue. The plug reinforces the soft tissue and then acts as a scaffold into which new tissue can grow to close the fistula. The plug is usually absorbed into the body in 6 to 8 weeks. The procedure can be repeated in case of failure.

RATIONALE:

The SIS Fistula Plug from Cook Biotech received 510(k) clearance from the U.S. Food and Drug Administration (FDA) in March 2005 based on similarity to predicate devices, including the SURGISIS® Soft Tissue Graft and the STRATASIS® Urethral Sling, both manufactured by Cook Biotech Incorporated. The SIS Fistula Plug is manufactured from porcine small intestinal submucosa (SIS) and is intended for repair of anal, rectal, and enterocutaneous fistulas. The modified SIS Fistula Plug, also manufactured from porcine small intestinal submucosa (SIS), is supplied in a tapered configuration with a button to provide increased retention of the plug and improved blockage of the fistula. It received 510(k) clearance in October 2006. In March 2009, W.L. Gore & Associates received 510(k) clearance for the BIO-A® Fistula Plug intended for use in anorectal fistulas. The GORE BIO-A Fistula Plug device comprises a porous structure of synthetic bioabsorbable PGA/TMC copolymer fiber, degraded via a

SUBJECT: PLUGS FOR FISTULA REPAIR	EFFECTIVE DATE: 08/18/11
POLICY NUMBER: 7.01.86	REVISED DATE: 07/19/12, 06/20/13, 05/22/14
CATEGORY: Technology Assessment	PAGE: 2 OF: 6

combination of hydrolytic and enzymatic pathways, the same material, technology, and three-dimensional disk with tubes mesh design as the predicate GORE Bioabsorbable Mesh hernia plug device. The indications for use and performance of the GORE BIO-A™ Fistula Plug are substantially equivalent to the predicate Cook SIS Fistula Plug.

H Ortiz and colleagues (2009) in a European trial compared use of porcine submucosal (Surgisis) anal fistula plug (AFP) with an endorectal anal flap (ERAF) procedure in a randomized controlled trial (RCT) with 43 patients who had high anal fistula. The primary endpoint was fistula healing. Recurrence was defined as the presence of an abscess in the same area or obvious evidence of fistulization. Five patients in the AFP group and 6 in the ERAF group did not receive the allocated intervention, leaving 32 patients. One patient in the AFP group was lost to follow-up. A large number of recurrences in the fistula plug group led to premature closure of the trial. After 1 year, fistula recurrence was seen in 12 of 15 patients treated with an anal fistula plug versus 2 of 16 patients who underwent the flap procedure (relative risk 6.40 [95% confidence interval 1.70-23.97]); p less than 0.001). Fistulas recurred in 9 of 16 patients who had previously undergone fistula surgery; 8 of the 9 patients had an AFP. A trend for more sphincter involvement and more females in the ERAF group was noted. Complications were not reported in this paper.

PJ van Koperen, et al. (2008) conducted a RCT to compare a fistula plug (n=31) with a mucosal advancement flap (n=29) for the treatment of high transsphincteric fistulas. At a follow-up of 11 months, the recurrence rates were 71% (n=22) in the anal fistula plug group and 52% (n=15) in the mucosal advancement flap group, which was not significantly different (p=.126). There were no significant differences in postoperative pain, in pre- and postoperative incontinence scores, soiling, and quality of life. One patient in plug group and two in flap group experienced postoperative complications (abscess, pain, bleeding retrospectively).

D Christoforidis, et al. (2009) performed a retrospective analysis of patients from a U.S. center with transsphincteric fistulas treated with ERAF (n=43) or anal plug (Surgisis) (n=37) between January 1996 and April 2007. Success was defined as closed external opening in absence of symptoms at minimal follow-up of 6 months. The success rate was 63% in the ERAF group and 32% in the AFP group after a mean follow-up of 56 (range, 6-136) months for ERAF and 14 (range, 6-22) months. After exclusion of patients with early AFP extrusion, which may be considered a technical failure, the ERAF advantage did not meet statistical significance (p=0.06). Twenty-three of 27 patients who had ERAF and 7 of 12 patients who had AFP responded to a questionnaire addressing functional outcomes. In the ERAF group 11 of 23 patients had no continence disturbance versus 6 of 7 in the AFP group. The lack of prospectively collected incontinence scores prior to the procedure and low response rate in the AFP group prohibit valid comparisons on functional outcomes. Complication rates were low in both groups; 2 patients in the ERAF group required reoperation for bleeding. No serious complications occurred in the AFP group. The authors conclude that “randomized trials are needed to further elucidate the efficacy and potential functional benefit of AFP in the treatment of complex anal fistulas.”

Wang, et al. (2009) compared outcomes of all patients with transsphincteric fistulas treated with AFP from July 2005 to December 2006 (n=29) and compared them with historical controls treated with ERAF (2001-2005) (n=26). Of 26 initial flap procedures, 10 failed and 16 healed. Of 29 initial plug procedures, 19 failed and 10 healed. In total, 30 advancement flaps and 34 plug procedures were performed (including the additional treatments for failed initial procedures). Closure rates were 34% for plugs (mean follow-up 279 [range, 110-690] days) and 62% for flaps (median follow-up 819 [range, 93-1928] days; p=0.045). Complications were not reported. The authors conclude that a systematic randomized trial with long-term follow-up comparing advancement flaps with fistula plugs is needed, and they calculate that 112 patients would need to be randomized to detect a statistically significant difference in success rates for each procedure. Because the fistula plugs are costly, the authors recommend that a cost-benefit analysis be performed.

A 2009 systematic review to access the efficacy of the anal fistula plug by P Garg and colleagues reports a wide range of success rates. In the 12 included studies, all case series, reported success rates for the AFP procedure were from 24% to 92%. Success rates in treating complex fistula-in-ano in the 8 prospective studies reviewed were 35%-87%. The authors concluded that while the anal fistula plug procedure appeared safe, further RCTs are needed.

In 2012, 3 reviews were published comparing AFP to conventional surgical treatment for anal fistulas. Pu and colleagues undertook a meta-analysis of 5 studies (2 RCTs and 3 retrospective studies) published through April 2012. Treatment options in the conventional arm of this review included endorectal/mucosal advancement flaps, fibrin glue,

SUBJECT: PLUGS FOR FISTULA REPAIR	EFFECTIVE DATE: 08/18/11
POLICY NUMBER: 7.01.86	REVISED DATE: 07/19/12, 06/20/13, 05/22/14
CATEGORY: Technology Assessment	PAGE: 3 OF: 6

and seton drains. On combined analysis, AFP patients had a higher recurrence rate (62%) compared to those undergoing conventional treatment options (47%) after 3 months of follow-up (5 studies, 428 patients; $p=0.004$).

Leng and Jin undertook a meta-analysis of 6 studies published through April 2011 (3 RCTs, 2 retrospective studies, and one cohort study) involving 408 patients comparing AFP with mucosal advancement flap (MAF). On combined analysis, the differences in the overall success rates (6 studies) and incidence of fistula recurrence (4 studies including 3 RCTs) were not statistically significant between the AFP and MAF. The risk of continence postoperatively (3 studies including 2 RCTs), however, was reported to be lower with AFP. In addition to the small numbers of controlled studies and limited follow-up, the findings of this meta-analysis were further limited by significant heterogeneity across studies.

O’Riordan and colleagues undertook a systematic review of AFP for patients with Crohn’s and non-Crohn’s-related anal fistulas. The follow-up period across studies ranged from 3 months to 24.5 months. The pooled proportion of patients achieving fistula closure in patients with non-Crohn’s anal fistula was 0.54. The proportion achieving closure in patients with Crohn’s disease was similar. There were no reported cases of any significant change in continence after AFP insertion in any of the study patients ($n=196$). The findings of this systematic review are limited by the variability of operative technique and perioperative care across studies, which may influence the probability of success or failure associated with the AFP.

Overall, the evidence of efficacy of anal fistula plug treatment is limited. Two randomized controlled trials and retrospective comparisons did not demonstrate that anal plugs improved healing rates or reduced recurrence of anal fistulas. Numerous case series report a wide range of results (e.g., CN Ellis, et al. 2010, BJ Champagne, et al. 2006, MF McGee, et al. 2010, S Gonsalves, et al. 2009) and contribute to the inability to allow conclusions to be drawn related to the long-term efficacy of fistula plugs. Randomized controlled trials with sufficient numbers of patients and with appropriate length of follow-up reporting healing and recurrence rates, and sphincter function, before and after procedures, are required.

CODES: Number Description

Eligibility for reimbursement is based upon the benefits set forth in the member’s subscriber contract.

CODES MAY NOT BE COVERED UNDER ALL CIRCUMSTANCES. PLEASE READ THE POLICY AND GUIDELINES STATEMENTS CAREFULLY.

Codes may not be all inclusive as the AMA and CMS code updates may occur more frequently than policy updates.

Code Key: Experimental/Investigational = (E/I), Not medically necessary/ appropriate = (NMN).

CPT: 46707 (E/I) Repair of anorectal fistula with plug (e.g., porcine small intestine mucosa [SIS])

Copyright© 2014 American Medical Association, Chicago, IL

HCPCS: No specific codes

<u>ICD9:</u>	565.1 Anal fistula
	569.81 Fistula of intestine, excluding rectum and anus (includes enterocutaneous or colocutaneous fistula)
	619.1 Digestive-genital tract fistula, female (includes rectovaginal)
<u>ICD10:</u>	K50.013 Crohn's disease of small intestine with fistula
	K50.113 Crohn's disease of large intestine with fistula
	K50.813 Crohn's disease of both small and large intestine with fistula
	K50.913 Crohn's disease, unspecified, with fistula
	K51.013 Ulcerative (chronic) pancolitis with fistula
	K5.1213 Ulcerative (chronic) proctitis with fistula

SUBJECT: PLUGS FOR FISTULA REPAIR	EFFECTIVE DATE: 08/18/11
POLICY NUMBER: 7.01.86	REVISED DATE: 07/19/12, 06/20/13, 05/22/14
CATEGORY: Technology Assessment	PAGE: 4 OF: 6

K51.313	Ulcerative (chronic) rectosigmoiditis with fistula
K51.413	Inflammatory polyps of colon with fistula
K51.513	Left sided colitis with fistula
K51.813	Other ulcerative colitis with fistula
K51.913	Ulcerative colitis, unspecified with fistula
K60.3-K60.5	Rectal anal fistula (code range)
K63.2	Fistula of intestine
N82.2-N82.4	Female intestinal-genital tract fistula (code range)

REFERENCES:

- Adamina M, et al. To plug or not to plug: a cost effectiveness analysis for complex anal fistula. *Surgery* 2010 Jan;147(1):72-8.
- Adamina M, et al. Anal fistula plug: a prospective evaluation of success, continence, and quality of life in the treatment of complex fistulas. *Colorectal Dis* 2014 Feb 13 [Epub ahead of print].
- Blom J, et al. Results of collagen plug occlusion of anal fistula: a multicenter study of 126 patients. *Colorectal Dis* 2014 Feb 10 [Epub ahead of print].
- BlueCross BlueShield Association. Plug for Fistula Repair. Medical Policy Reference Manual. Policy #7.01.123. 2013 Sep 12.
- Champagne BJ, et al. Rectovaginal fistula. *Surg Clin North Am* 2010 Feb;90(1):69-82.
- *Champagne BJ, et al. Efficacy of anal fistula plug in closure of cryptoglandular fistulas: long-term follow-up. *Dis Colon Rectum* 2006 Dec;49(12):1817-21.
- Chan S, et al. Initial experience of treating anal fistula with Surgisis anal fistula plug. *Tech Coloproctol* 2012 Jun;16(3):201-6.
- *Christoforidis D, et al. Treatment of transsphincteric anal fistulas by endorectal advancement flap or collagen fistula plug: a comparative study. *Dis Colon Rectum* 2009 Jan;52(1):18-22.
- Cintron JR, et al. Treatment of fistula-in-ano using a porcine small intestinal submucosa anal fistula plug. *Tech Coloproctol* 2013 Apr;17(2):187-91.
- Dela Portilla F, et al. Evaluation of a new synthetic plug in the treatment of anal fistulas: results of a pilot study. *Dis Colon Rectum* 2011 Nov;54(11):1419-22.
- El-Gazzaz G, et al. A retrospective review of chronic anal fistulae treated by anal fistulae plug. *Colorectal Dis* 2010 May;12(5):442-7.
- Ellis CN, et al. Long-term outcomes with the use of bioprosthetic plugs for the management of complex anal fistulas. *Dis Colon Rectum* 2010 May;53(5):798-802.
- Gajsek U, et al. Long-term efficacy of the button fistula plug in the treatment of ileal pouch-vaginal and Crohn's-related rectovaginal fistulas. *Dis Colon Rectum* 2011 Aug;54(8):999-1002.
- *Garg P, et al. The efficacy of anal fistula plug in fistula-in-ano: a systematic review. *Colorectal Dis* 2010 Oct;12(10):965-70.
- Gonsalves S, et al. Assessment of efficacy of the rectovaginal button fistula plug for the treatment of ileo pouch-vaginal and rectovaginal fistulas. *Dis Colon Rectum* 2009 Nov;52(11):1877-81.
- Han JG, et al. Long-term outcomes of human acellular dermal matrix plug in closure of complex anal fistulas with a single tract. *Dis Colon Rectum* 2011 Nov;54(11):1412-8.

SUBJECT: PLUGS FOR FISTULA REPAIR	EFFECTIVE DATE: 08/18/11
POLICY NUMBER: 7.01.86	REVISED DATE: 07/19/12, 06/20/13, 05/22/14
CATEGORY: Technology Assessment	PAGE: 5 OF: 6

- Han JG, et al. Ligation of the intersphincteric fistula tract plus a bioprosthetic anal fistula plug (LIFT-Plug): a new technique for fistula-in-ano. Colorectal Dis 2013 May;15(5):582-6.
- Heydari A, et al. Bioabsorbable synthetic plug in the treatment of anal fistulas. Dis Colon Rectum 2013 Jun;56(6):774-9.
- Hyman N, et al. Outcomes after fistulotomy: results of a prospective, multicenter regional study. Dis Colon Rectum 2009 Dec;52(12):2022-7.
- Jacob TJ, et al. Surgical intervention for anorectal fistula. Cochrane Database Syst Rev 2010 May 12;(5):CD006319.
- *Johnson EK, et al. Efficacy of anal fistula plug vs fibrin glue in closure of anorectal fistulas. Dis Colon Rectum 2006 Mar;49(3):371-6.
- Leng Q, et al. Anal fistula plug vs mucosa advancement flap in complex fistula-in-ano: A met-analysis. World J Gastrointest Surg 2012 Nov 27;4(11):256-61.
- Lenisa L, et al. Anal fistula plug is a valid alternative option for the treatment of complex anal fistula in the long term. Int J Colorectal Dis 2010 Dec;25(12):1487-93.
- Lupinacci RM, et al. Treatment of fistula-in-ano with the Surgisis® AFP™ anal fistula plug. Gastroenterol Clin Biol 2010 Oct;34(10):549-53.
- McGee MF, et al. Tract length predicts successful closure with anal fistula plug in cryptoglandular fistulas. Dis Colon Rectum 2010 Aug;53(8):1116-20.
- *National Institute for Health and Excellence (NICE). Closure of anal fistula using a suturable bioprosthetic plug. 2012 May [http://www.nice.org.uk/guidance/index.jsp] accessed 4/17/14.
- Pinto RA, et al. Are there predictors of outcome following rectovaginal fistula repair? Dis Colon Rectum 2010 Sep;53(9):1240-7.
- *O'Connor L, et al. Efficacy of anal fistula plug in closure of Crohn's anorectal fistulas. Dis Colon Rectum 2006 Oct;49(10):1569-73.
- Ommer A, et al. Gore BioA fistula plug in the treatment of high anal fistulas- initial results from a German multi-center study. Ger Med Sci 2012;10:Doc13.
- O'Riordan JM, et al. A systematic review of the anal fistula plug for patients with Crohn's and non-Crohn's related fistula-in-ano. Dis Colon Rectum 2012 Mar;55(3):351-8.
- *Ortiz H, et al. Randomized clinical trial of anal fistula plug versus endorectal advancement flap for the treatment of high cryptoglandular fistula in ano. Br J Surg 2009 Jun;96(6):608-12.
- Pu YW, et al. Fistula plug versus conventional surgical treatment for anal fistulas. A system review and meta-analysis. Saudi Med J 2012 Sep;33(9):962-6.
- Ratto C, et al. Gore Bio-A® Fistula Plug; a new sphincter-sparing procedure for complex anal fistula. Colorectal Dis 2012 May;14(5):e264-9.
- Rizzo JA, et al. Anorectal abscess and fistula-in-ano: evidence-based management. Surg Clin North AM 2010 Feb;90(1):45-68.
- Safar B, et al. Anal fistula plug: initial experience and outcomes. Dis Colon Rectum 2009 Feb;52(2):248-52.
- *Schwandner O, et al. Initial experience on efficacy in closure of cryptoglandular and Crohn's transsphincteric fistulas by the use of the anal fistula plug. In J Colorectal Dis 2008 Mar;23(3):319-24.
- Schwandner O, et al. Preliminary results on efficacy in closure of transsphincteric and rectovaginal fistulas associated with Crohn's disease using new biomaterials. Surg Innov 2009 Jun;16(2):162-8.
- Schwandner T, et al. Surgical treatment of complex anal fistulas with the anal fistula plug: a prospective, multicenter study. Dis Colon Rectum 2009 Sep;52(9):1578-83.

SUBJECT: PLUGS FOR FISTULA REPAIR	EFFECTIVE DATE: 08/18/11
POLICY NUMBER: 7.01.86	REVISED DATE: 07/19/12, 06/20/13, 05/22/14
CATEGORY: Technology Assessment	PAGE: 6 OF: 6

Tan KK, et al. Long-term outcome of the anal fistula plug for anal fistula of cryptoglandular origin. *Colorectal Dis* 2013 Dec;15(12):1510-4.

Thekkinkattil DK, et al. Efficacy of the anal fistula plug in complex anorectal fistulae. *Colorectal Dis* 2009 Jul;11(6):584-7.

*Van Koperen PJ, et al. Anal fistula plug for closure of difficult anorectal fistula: a prospective study. *Dis Colon Rectum* 2007 Dec;50(12):2168-72.

*Van Koperen PJ, et al. The anal fistula plug treatment compared with mucosal advancement flap for cryptoglandular high transsphincteric perianal fistula: a double-blinded multicenter randomized trial. *BMC Surg* 2008 Jun 23;8:11.

Van Koperen PJ, et al. The anal fistula plug treatment compared with mucosal advancement flap for cryptoglandular high transsphincteric perianal fistula: a double-blinded multicenter randomized trial. *Dis Colon Rectum* 2011 Apr;54(4):387-93.

Wang JY, et al. Treatment of transsphincteric anal fistulas: are fistula plugs an acceptable alternative? *Dis Colon Rectum* 2009 Apr;52(4):692-7.

*Whiteford MH, et al. Practice parameters for the treatment of perianal abscess and fistula-in-ano (revised). *Dis Colon Rectum* 2005 Jul;48(7):1337-42.

* key article

KEY WORDS:

Fistula plug

CMS COVERAGE FOR MEDICARE PRODUCT MEMBERS

Based upon our review, repair of an anal fistula with a fistula plug is not addressed in National or regional CMS coverage determinations or policies.