

Protocol

Intraoperative Neurophysiologic Monitoring (sensory-evoked potentials, motor-evoked potentials, EEG monitoring)

(70158)

Medical Benefit	Effective Date: 07/01/14	Next Review Date: 05/15
Preauthorization	No	Review Dates: 03/13, 05/13, 05/14

*The following Protocol contains medical necessity criteria that apply for this service. It is applicable to Medicare Advantage products unless separate Medicare Advantage criteria are indicated. If the criteria are not met, reimbursement will be denied and the patient cannot be billed. **Preauthorization is not required.** Please note that payment for covered services is subject to eligibility and the limitations noted in the patient's contract at the time the services are rendered.*

Description

Intraoperative neurophysiologic monitoring (IONM) describes a variety of procedures that have been used to monitor the integrity of neural pathways during high-risk neurosurgical, orthopedic, and vascular surgeries. It involves the detection of electrical signals produced by the nervous system in response to sensory or electrical stimuli to provide information about the functional integrity of neuronal structures.

Background

The principal goal of intraoperative neurophysiologic monitoring (IONM) is the identification of nervous system impairment in the hope that prompt intervention will prevent permanent deficits. Correctable factors at surgery include circulatory disturbance, excess compression from retraction, bony structures, or hematomas, or mechanical stretching. The technology is continuously evolving with refinements in equipment and analytic techniques, including recording, with several patients monitored under the supervision of a physician who is outside the operating room.

The different methodologies of monitoring are described below:

Sensory-evoked Potentials

Sensory-evoked potential describes the responses of the sensory pathways to sensory or electrical stimuli. Intraoperative monitoring of sensory-evoked potentials is used to assess the functional integrity of central nervous system (CNS) pathways during operations that put the spinal cord or brain at risk for significant ischemia or traumatic injury. The basic principles of sensory-evoked potential monitoring involve identification of a neurological region at risk, selection and stimulation of a nerve that carries a signal through the at-risk region, and recording and interpretation of the signal at certain standardized points along the pathway. Monitoring of sensory-evoked potentials is commonly used during the following procedures: carotid endarterectomy, brain surgery involving vasculature, surgery with distraction compression or ischemia of the spinal cord and brainstem, and acoustic neuroma surgery. Sensory-evoked potentials can be further broken down into the following categories according to the type of simulation used:

- Somatosensory-evoked potentials (SSEPs) are cortical responses elicited by peripheral nerve stimulations. Peripheral nerves, such as the median, ulnar, or tibial nerves, are typically stimulated, but in some situations the spinal cord may be stimulated directly. Recording is done either cortically or at the level of the spinal cord above the surgical procedure. Intraoperative monitoring of SSEPs is most commonly used during orthopedic or neurologic surgery to prompt intervention to reduce surgically induced morbidity and/or to monitor the level of anesthesia. One of the most common indications for SSEP monitoring is in patients

undergoing corrective surgery for scoliosis. In this setting, SSEP monitors the status of the posterior column pathways and thus does not reflect ischemia in the anterior (motor) pathways. Several different techniques are commonly used, including stimulation of a relevant peripheral nerve with monitoring from the scalp, from interspinous ligament needle electrodes, or from catheter electrodes in the epidural space.

- Brainstem auditory-evoked potentials (BAEPs) are generated in response to auditory clicks and can define the functional status of the auditory nerve. Surgical resection of a cerebellopontine angle tumor, such as an acoustic neuroma, places the auditory nerves at risk, and BAEPs have been extensively used to monitor auditory function during these procedures.
- Visual-evoked potentials (VEPs) with light flashes are used to track visual signals from the retina to the occipital cortex. VEP monitoring has been used for surgery on lesions near the optic chiasm. However, VEPs are very difficult to interpret due to their sensitivity to anesthesia, temperature, and blood pressure.

Motor-evoked Potentials

Motor-evoked potentials (MEPs) are recorded from muscles following direct or transcranial electrical stimulation of motor cortex or by pulsed magnetic stimulation provided by a coil placed over the head. Peripheral motor responses (muscle activity) are recorded by electrodes placed on the skin at prescribed points along the motor pathways. Motor evoked potentials, especially when induced by magnetic stimulation, can be affected by anesthesia. The Digitimer electrical cortical stimulator received U.S. Food and Drug Administration (FDA) premarket approval in 2002. Devices for transcranial magnetic stimulation have not yet received approval from the FDA for this use.

Multimodal IONM, in which more than one technique is used, most commonly with SSEPs and MEPs, has also been described.

EMG (Electromyogram) Monitoring and Nerve Conduction Velocity Measurements

Electromyogram monitoring and nerve conduction velocity measurements can be performed in the operating room and may be used to assess the status of the peripheral nerves, e.g., to identify the extent of nerve damage prior to nerve grafting or during resection of tumors. In addition, these techniques may be used during procedures around the nerve roots and around peripheral nerves to assess the presence of excessive traction or other impairment. Surgery in the region of cranial nerves can be monitored by electrically stimulating the proximal (brain) end of the nerve and recording via EMG in the facial or neck muscles. Thus monitoring is done in the direction opposite that of sensory-evoked potentials, but the purpose is similar—to verify that the neural pathway is intact.

EEG (Electroencephalogram) Monitoring

Spontaneous EEG monitoring can also be recorded during surgery and can be subdivided as follows:

- EEG monitoring has been widely used to monitor cerebral ischemia secondary to carotid cross-clamping during a carotid endarterectomy. EEG monitoring may identify those patients who would benefit from the use of a vascular shunt during the procedure to restore adequate cerebral perfusion. Conversely, shunts, which have an associated risk of iatrogenic complications, may be avoided in those patients in whom the EEG is normal. Carotid endarterectomy may be done with the patient under local anesthesia so that monitoring of cortical function can be directly assessed.
- Electrocorticography (ECoG) is the recording of the EEG directly from a surgically exposed cerebral cortex. CoG is typically used to define the sensory cortex and map the critical limits of a surgical resection. ECoG recordings have been most frequently used to identify epileptogenic regions for resection. In these applications, ECoG does not constitute monitoring, per se.

Policy (Formerly Corporate Medical Guideline)

Intraoperative monitoring, which includes somatosensory-evoked potentials, motor-evoked potentials using transcranial electrical stimulation, brainstem auditory-evoked potentials, EMG of cranial nerves, EEG, and electrocorticography (ECoG), may be considered **medically necessary** during spinal, intracranial, or vascular procedures.

Intraoperative monitoring of visual-evoked potentials is considered **investigational**.

Due to the lack of FDA approval, intraoperative monitoring of motor-evoked potentials using transcranial magnetic stimulation is considered **investigational**.

Intraoperative EMG and nerve conduction velocity monitoring during surgery on the peripheral nerves is considered **not medically necessary**.

Note: These policy statements refer only to use of these techniques as part of intraoperative monitoring. Other clinical applications of these techniques, such as visual-evoked potentials and EMG, are not considered in this Protocol.

Policy Guideline

Constant communication between surgeon, neurophysiologist, and anesthetist are required for safe and effective intraoperative neurophysiologic monitoring.

Medicare Advantage

In addition to the above, EEG monitoring during open heart surgery would be considered **not medically necessary** for Medicare Advantage members.

Benefit Application

Intraoperative monitoring is considered reimbursable as a separate service only when a licensed healthcare practitioner, other than the operating surgeon, interprets the monitoring. The provision/monitoring of the sensory procedure is performed by a healthcare practitioner or technician who is in attendance in the operating room throughout the procedure. The intraoperative monitoring performed remotely would be provided in conjunction with the healthcare provider who is providing/monitoring the sensory test in the operating room, by the remote physician whose attention is directed solely on one patient.

Those providing the intraoperative monitoring services must have the proper training to do so whether providing real-time review and interpretation on-site versus remote.

Services that are the subject of a clinical trial do not meet our Technology Assessment Protocol criteria and are considered investigational. *For explanation of experimental and investigational, please refer to the Technology Assessment Protocol.*

It is expected that only appropriate and medically necessary services will be rendered. We reserve the right to conduct prepayment and postpayment reviews to assess the medical appropriateness of the above-referenced

procedures. **Some of this Protocol may not pertain to the patients you provide care to, as it may relate to products that are not available in your geographic area.**

References

We are not responsible for the continuing viability of web site addresses that may be listed in any references below.

1. Aminoff MJ. Intraoperative monitoring by evoked potentials for spinal cord surgery: the cons. *Electroencephalogr Clin Neurophysiol* 1989; 73(5):378-80.
2. Daube JR. Intraoperative monitoring by evoked potentials for spinal cord surgery: the pros. *Electroencephalogr Clin Neurophysiol* 1989; 73(5):374-7.
3. Fisher RS, Raudzens P, Nunemacher M. Efficacy of intraoperative neurophysiological monitoring. *J Clin Neurophysiol* 1995; 12(1):97-109.
4. Schweiger H, Kamp HD, Dinkel M. Somatosensory-evoked potentials during carotid artery surgery: experience in 400 operations. *Surgery* 1991; 109(5):602-9.
5. Fehlings MG, Brodke DS, Norvell DC et al. The evidence for intraoperative neurophysiological monitoring in spine surgery: does it make a difference? *Spine (Phila Pa 1976)* 2010; 35(9 Suppl):S37-46.
6. Nuwer MR, Emerson RG, Galloway G et al. Evidence-based guideline update: intraoperative spinal monitoring with somatosensory and transcranial electrical motor evoked potentials: report of the Therapeutics and Technology Assessment Subcommittee of the American Academy of Neurology and the American Clinical Neurophysiology Society. *Neurology* 2012; 78(8):585-9.
7. Schwartz DM, Auerbach JD, Dormans JP et al. Neurophysiological detection of impending spinal cord injury during scoliosis surgery. *J Bone Joint Surg Am* 2007; 89(11):2440-9.
8. Ota T, Kawai K, Kamada K et al. Intraoperative monitoring of cortically recorded visual response for posterior visual pathway. *J Neurosurg* 2010; 112(2):285-94.
9. Sasaki T, Itakura T, Suzuki K et al. Intraoperative monitoring of visual evoked potential: introduction of a clinically useful method. *J Neurosurg* 2010; 112(2):273-84.
10. Malhotra NR, Shaffrey CI. Intraoperative electrophysiological monitoring in spine surgery. *Spine (Phila Pa 1976)* 2010; 35(25):2167-79.
11. American Electroencephalographic Society. Guidelines for clinical evoked potential studies. *J Clin Neurophysiol* 1984; 1(1):3-53.
12. American Electroencephalographic Society. Guideline eleven: guidelines for intraoperative monitoring of sensory evoked potentials. *J Clin Neurophysiol* 1994; 11(1):77-87.
13. American Clinical Neurophysiology Society. Guideline 11A. Recommended Standards for Neurophysiologic Intraoperative Monitoring – Principles. Available online at: <http://www.acns.org/pdfs/11A%20-%20Recommended%20Standards%20for%20NIOM%20-%20Principles.pdf>. Last accessed November, 2012.
14. American Academy of Neurology. Assessment: intraoperative neurophysiology. Report of the Therapeutics and Technology Assessment Subcommittee of the American Academy of Neurology. *Neurology* 1990; 40(11):1644-6.
15. American Society of Neurophysiologic Monitoring. Position Statements. Available online at: <http://www.asnm.org/news/position-statements.html>. Last accessed November, 2012.

16. International Organization of Societies for Electrophysiological Technology (OSET). Guidelines for Performing EEG and Evoked Potential Monitoring During Surgery. Am J END Technol 1999; 39:257-77.
17. Nuwer MR, Daube J, Fischer C et al. Neuromonitoring during surgery. Report of an IFCN Committee. Electroencephalogr Clin Neurophysiol 1993; 87(5):263-76.
18. Centers for Medicare and Medicaid Services. National Coverage Determination (NCD) for Electroencephalographic monitoring During Surgical Procedures Involving the Cerebral Vasculature (160.8). Available online at: <http://www.cms.gov/medicare-coverage-database/details/ncd-details.aspx?NCDId=77&ncdver=2&CoverageSelection=National&Keyword=monitoring&KeywordLookUp=Title&KeywordLookUp=Title&KeywordLookUp=Title&KeywordSearchType=And&KeywordSearchType=And&KeywordSearchType=And&bc=gAAAAACAAAAA&>. Last accessed April, 2014.
19. Centers for Medicare and Medicaid Services National Coverage Determination (NCD) for Electroencephalographic (EEG) Monitoring During Open-Heart Surgery (160.9), Effective Date of this Version. This is a longstanding national coverage determination. The effective date of this version has not been posted.