

Protocol

Small Bowel/Liver and Multivisceral Transplant

(70305)

| | | |
|------------------|--------------------------|---|
| Medical Benefit | Effective Date: 07/01/14 | Next Review Date: 05/15 |
| Preauthorization | Yes | Review Dates: 05/09, 05/10, 05/11, 05/12, 05/13, 07/13, 05/14 |

*The following Protocol contains medical necessity criteria that apply for this service. It is applicable to Medicare Advantage products unless separate Medicare Advantage criteria are indicated. If the criteria are not met, reimbursement will be denied and the patient cannot be billed. **Preauthorization is required and must be obtained through Case Management.** Please note that payment for covered services is subject to eligibility and the limitations noted in the patient's contract at the time the services are rendered.*

Description

Small bowel/liver transplantation is transplantation of an intestinal allograft in combination with a liver allograft, either alone or in combination with one or more of the following organs: stomach, duodenum, jejunum, ileum, pancreas, or colon.

Background

Small bowel transplants are typically performed in patients with short bowel syndrome, defined as an inadequate absorbing surface of the small intestine due to extensive disease or surgical removal of a large portion of small intestine. In some instances, short bowel syndrome is associated with liver failure, often due to the long-term complications of total parenteral nutrition (TPN). These patients may be candidates for a small bowel/liver transplant or a multivisceral transplant, which includes the small bowel and liver with one or more of the following organs: stomach, duodenum, jejunum, ileum, pancreas, and/or colon. A multivisceral transplant is indicated when anatomic or other medical problems preclude a small bowel/liver transplant.

Related Protocol

Isolated Small Bowel Transplant

Policy (Formerly Corporate Medical Guideline)

A small bowel/liver transplant or multivisceral transplant may be considered **medically necessary** for pediatric and adult patients with intestinal failure (characterized by loss of absorption and the inability to maintain protein-energy, fluid, electrolyte, or micronutrient balance), who have been managed with long-term total parenteral nutrition (TPN) and who have developed evidence of impending end-stage liver failure.

A small bowel/liver retransplant or multivisceral retransplant may be considered **medically necessary** after a failed primary small bowel/liver transplant or multivisceral transplant.

Small bowel/liver transplant or multivisceral transplant would be considered **investigational** in all other situations.

Policy Guideline

General

Potential contraindications to solid organ transplant (subject to the judgment of the transplant center):

1. Known current malignancy, including metastatic cancer
2. Recent malignancy with high risk of recurrence
3. History of cancer with a moderate risk of recurrence
4. Systemic disease that could be exacerbated by immunosuppression
5. Untreated systemic infection making immunosuppression unsafe, including chronic infection
6. Other irreversible end-stage disease not attributed to intestinal failure
7. Psychosocial conditions or chemical dependency affecting ability to adhere to therapy.

Intestinal failure results from surgical resection, congenital defect, or disease-associated loss of absorption and is characterized by the inability to maintain protein-energy, fluid, electrolyte, or micronutrient balance (adapted from reference 1). (1) Short-bowel syndrome is one case of intestinal failure.

Candidates should meet the following criteria:

- Adequate cardiopulmonary status
- Documentation of patient compliance with medical management.

HIV [human immunodeficiency virus]-positive patients who meet the following criteria, as stated in the 2001 guidelines of the American Society of Transplantation, could be considered candidates for small bowel/liver or multivisceral transplantation:

- CD4 count greater than 200 cells per cubic millimeter for greater than six months
- HIV-1 RNA undetectable
- On stable anti-retroviral therapy greater than three months
- No other complications from AIDS [acquired immune deficiency syndrome] (e.g., opportunistic infection, including aspergillus, tuberculosis, coccidiosis mycosis, resistant fungal infections, Kaposi's sarcoma, or other neoplasm), and meeting all other criteria for transplantation.

Small Bowel/Liver Specific

Evidence of intolerance of total parenteral nutrition (TPN) includes, but is not limited to, multiple and prolonged hospitalizations to treat TPN-related complications, or the development of progressive but reversible liver failure. In the setting of progressive liver failure, small bowel transplant may be considered a technique to avoid end-stage liver failure related to chronic TPN, thus avoiding the necessity of a multivisceral transplant.

Benefit Application

Individual transplant facilities may have their own *additional* requirements or protocols that must be met in order for the patient to be eligible for a transplant at **their** facility.

Medicare Advantage

If a transplant is needed, we arrange to have the Medicare-approved transplant center review and decide whether the patient is an appropriate candidate for the transplant.

Services that are the subject of a clinical trial do not meet our Technology Assessment Protocol criteria and are considered investigational. *For explanation of experimental and investigational, please refer to the Technology Assessment Protocol.*

It is expected that only appropriate and medically necessary services will be rendered. We reserve the right to conduct prepayment and postpayment reviews to assess the medical appropriateness of the above-referenced procedures. **Some of this Protocol may not pertain to the patients you provide care to, as it may relate to products that are not available in your geographic area.**

References

We are not responsible for the continuing viability of web site addresses that may be listed in any references below.

1. O'Keefe SJ, Buchman AL, Fishbein TM et al. Short bowel syndrome and intestinal failure: consensus definitions and overview. Clin Gastroenterol Hepatol 2006; 4(1):6-10.
2. Blue Cross and Blue Shield Association Technology Evaluation Center (TEC). Small bowel transplants in adults and multivisceral transplants in adults and children. TEC Assessments 1999; Volume 14, Tab 9.
3. Nayyar N, Mazariegos G, Ranganathan S et al. Pediatric small bowel transplantation. Semin Pediatr Surg 2010; 19(1):68-77.
4. Abu-Elmagd KM, Costa G, Bond GJ et al. Five hundred intestinal and multivisceral transplantations at a single center: major advances with new challenges. Ann Surg 2009; 250(4):567-81.
5. Varkey J, Simren M, Bosaeus I et al. Survival of patients evaluated for intestinal and multivisceral transplantation - the Scandinavian experience. Scand J Gastroenterol 2013; 48(6):702-11.
6. Mangus RS, Tector AJ, Kubal CA et al. Multivisceral transplantation: expanding indications and improving outcomes. J Gastrointest Surg 2013; 17(1):179-86; discussion p 86-7.
7. Wu G, Selvaggi G, Nishida S et al. Graft-versus-host disease after intestinal and multivisceral transplantation. Transplantation 2011; 91(2):219-24.
8. Florescu DF, Qiu F, Langnas AN et al. Bloodstream Infections during the First Year after Pediatric Small Bowel Transplantation. Pediatr Infect Dis J 2012; In Press.
9. Steinman TI, Becker BN, Frost AE et al. Guidelines for the referral and management of patients eligible for solid organ transplantation. Transplantation 2001; 71(9):1189-204.
10. Bhagani S, Sweny P, Brook G. Guidelines for kidney transplantation in patients with HIV disease. HIV Med 2006; 7(3):133-9.
11. United Network for Organ Sharing. Identification of transmissible diseases in organ recipients. Available online at: optn.transplant.hrsa.gov/PoliciesandBylaws2/policies/pdfs/policy_16.pdf. Last accessed April, 2012.
12. Trevizol AP, David AI, Yamashita ET et al. Intestinal and multivisceral retransplantation results: literature review. Transplant Proc 2013; 45(3):1133-6.
13. Mazariegos GV, Soltys K, Bond G et al. Pediatric intestinal retransplantation: techniques, management, and outcomes. Transplantation 2008; 86(12):1777-82.
14. Desai CS, Khan KM, Gruessner AC et al. Intestinal retransplantation: analysis of Organ Procurement and Transplantation Network database. Transplantation 2012; 93(1):120-5.

15. Center for Medicare and Medicaid Services. Intestinal and multi-visceral transplantation. Available online at: www.cms.gov. Last accessed May, 2013.